THOMASAW

86

85g - 180 pages

GIBERT JOSEPH

Cahiers Thomason - numérisé par la Bibliothèque MIR (Université Paris Diderot : Université Pierre et Marie Curie ; CNRS)

p 2 strongly left enriched

p 3 EQUIMORPHISM = weak equivalence (8 Sept 1995)

p 4 if $K \in \mathcal{K}_{cof}$ $L \rightarrowtail L'$ $\Rightarrow$ $K \otimes L \rightarrowtail K \otimes L'$
strong left enrichment restricts to a left enrichment

p 7 dense $\mathcal{C}_{cof} \subset \mathcal{C}$ need not preserve finite colimits!
eg. free fg $\mathbb{Z}$-mod $= \mathcal{P}$ has all finite colimits,
but $\mathcal{P} \subset \mathbb{Z}$-mod doesn't preserve them
So $\mathcal{C}^s_{cof}(\mathbb{Z}) \subset \mathcal{C}(\mathbb{Z})$ doesn't preserve all finite colimits

p 9: But $\mathcal{C}_{cof} \subset \mathcal{C}$ does preserve cobase change along cofibrations

p 13 Rich hypotheses: enriched over Δ^{op}-sets

p 15 Rich hypotheses self-dual

p 16 Prop: Rich Hypo $\Rightarrow$ $\otimes$ respects colimits,
$\triangleleft$ (colimit) $\times$ limit $\longrightarrow$ limit.

p 24 Rich Hypo $\Rightarrow$ $C \rightarrowtail C'$, $D' \twoheadrightarrow D$ $\quad \underline{Hom}(C', D') \longrightarrow \underline{Hom}(C, D') \times_{\underline{Hom}(C, D)} \underline{Hom}(C', D)$
and $\xrightarrow{\sim}$ if $C \xrightarrow{\sim} C'$ or $D' \xrightarrow{\sim} D$

p 28 Rich Hypo + $\mathcal{N}$ (product) + ω summable
$\Rightarrow$ weak hyp + strong hypo + extra hypo

p 38 $\underline{Hom}(C, \ )$ preserves limits, $\underline{Hom}(\ , C)$ takes colimits to limits

p 40 If $\mathcal{C}$ $\mathcal{N}$-complete, strongly $\mathcal{N}$-reproductive, ultra ω-cosummable
hocolim$_{\Delta/K}$ F exists for each $F: \Delta/K \longrightarrow \mathcal{C}$ (not in $\mathcal{C}_{cof}$ necessarily)

p 44 def of ultra ω-cosummable

p 49 40 holds for Δ^{op}-sets.

p 51 left model, left functorial fact, + enriched over Δ^{op}-sets $\Rightarrow$ $\otimes, D \Rightarrow$
weak, strong + extra hypotheses of §1, §3, §5

p 54 very generally $\underline{Hom}(\text{hocolim}_{\Delta/K} F, C) = \text{holim}_{(\Delta/K)^{op}} \underline{Hom}(F, C)$

$\int^{K} \underline{Hom}(\text{hocolim}_{K/K} FK, GK') \simeq \text{hoend}(K, K' \rightarrow \underline{Hom}_{\mathcal{C}}(FK, GK'))$

hocolim$_{\Delta/K}$ preserves colimits which exist in $\mathcal{C}$ & stay in $\mathcal{C}_{cof}$.
model

p 61. Enriched lift of $\mathcal{N}$ structure from $\mathcal{C}$ to $Cat(K, \mathcal{C})$!
with weak (all needed hypotheses on $\mathcal{C}$ lift to $Cat(K, \mathcal{C})$!)

Cahiers Thomason. Bibliothèque MIR

so

$$\amalg C \xrightarrow{\nabla} C$$

$$\operatorname{hocolim}\Big(\operatorname{hocolim} I'I'C \sqcup_{\operatorname{hocolim} I'(C\sqcup C)} I(C\sqcup C)\Big) \xrightarrow{\operatorname{hocolim}(h\to I\circ h)} \operatorname{hocolim}\Big(I'I'C \sqcup_{I'C\sqcup C} I(C\sqcup C)\Big)$$

$\simeq$? $\simeq$? $\simeq$?

$\uparrow$ "$I(i_0 \sqcup i_1)$" $\simeq$ $\uparrow$ $I\nabla\circ\sigma$.

$$\operatorname{hocolim} I'(C\sqcup C) \xrightarrow{\quad} \operatorname{hocolim} I'C\sqcup C$$

$\simeq$?

might possibly work. But will be a mess.

Another idea: If $\mathcal{C}$ left model category with functorial factorization, and $\mathcal{C}$ left enriched over Δ_f,

then $\exists$ some $\Delta_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$.

Set $K\tilde{\otimes}C$ to be $\emptyset \to \tilde{C} \xrightarrow{\sim} C$ functorial.

$K\tilde{\otimes}C =$ ~~pushout~~

$K\otimes\tilde{C}$

cf 80 r 3
reason for making
left invt
New invariant

1 st $K\otimes\tilde{C} \to \Delta[0]\otimes\tilde{C} \xrightarrow{\sim} C$

$\Delta[0]\tilde{\otimes}C = \Delta[0]\otimes\tilde{C} \xrightarrow{\sim} \tilde{C} \to C.$

Will work if The functorial factor $(\emptyset \to C) \mapsto \tilde{C}$

is st $C \hookrightarrow C' \Rightarrow \tilde{C} \hookrightarrow \tilde{C}'$ and $\sim$ preserves

pushouts along cofibration.

ie want
axioms
C1, C2, C3 in
cf 79 r 72

I was reluctant to assume

$\Delta_+ \otimes C \to C$ for all C

because Quillen says (Homotopical Algebra II p 0.2) that certain categories of DGA's don't seem to have a simplicial model structure

Cahiers Thomason. Bibliothèque MIR

I've skimmed through Rational Homotopy and "On the (Co-)homology of commutative rings" without seeing the problem discussed.

Hom enriched shows up in Rational Homotopy II 54, seemingly with any previous reference in the paper. But it is in category of simplicial objects sC.

—

Second reason for being reluctant to assume $1_f \otimes C \xrightarrow{\otimes} C$ was idea to try to build ~~a structure~~ such a structure from ~~the~~ functorial factorization, assuming some ~~I² → I~~ .. extra axioms. Eg. on 79, 120, 139-141, 159.

Cf. discussion of Problem on 80, p 3

But the approach to constructing an enrichment is now dormant.

~~Idea~~ Anyway, if natural preserves cofibrations and pushouts, etc, we can use idea of p 1 to extend

$1_f \times C_{cof} \xrightarrow{\otimes} C_{cof}$ so constructed (if possible) to

$1_f \times C \longrightarrow C$.

—

So Let's go for

Def.

~~Strongly left enriched over~~

Def. Strongly left enriched (cf p 6 (cf 81, p 51),

Let $\mathcal{K}, \mathcal{L}$ be two left semi-model categories (81, 48). A strong left enrichment of $\mathcal{L}$ over $\mathcal{K}$ is a functor $\mathcal{K} \times \mathcal{L} \xrightarrow{\otimes} \mathcal{L}$ such that

Cahiers Thomason. Bibliothèque MIR

LS
~~SK~~ I a 1) $\forall K \in \mathscr{K}$ $K \otimes : \mathcal{Z} \longrightarrow \mathcal{Z}$ sends $\emptyset$ to $\emptyset$ and sends a pushout square along a cofibration in $\mathcal{Z}$ to a pushout square in $\mathcal{Z}$. (I do not ask $K \otimes$ preserves cofibrations in general)

LS I a 2) $\forall L \in \mathcal{Z}$, $\otimes L : \mathscr{K} \longrightarrow \mathcal{Z}$ sends $\emptyset$ to $\emptyset$, and sends pushout squares ~~from~~ along a cofibration to pushout squares in $\mathcal{Z}$. (Do not ask $\otimes L$ preserves cofibrations in general)

LS I 5) EMPTY.

LS I c) If $L \rightarrowtail L'$ is a cofibration in $\mathcal{Z}$ and $K \rightarrowtail K'$ is a cofibration in $\mathscr{K}$, then the pushout

$$K' \otimes L \underset{K \otimes L}{\cup} K \otimes L'$$

is representable in $\mathcal{Z}$, and the induced map

$$K' \otimes L \underset{K \otimes L}{\cup} K \otimes L' \longrightarrow K' \otimes L'$$

is a cofibration.

Moreover, if either $L \rightarrowtail L'$ or $K \rightarrowtail K'$ is also a weak equivalence, then the induced map

$$K' \otimes L \underset{K \otimes L}{\cup} K \otimes L' \longrightarrow K' \otimes L'$$

is a weak equivalence.

Note: Examples show that one should not ask that ~~$K \otimes$~~ $\otimes L$ preserve cofibrations if L not cofibrant. (cf §0 n° 3).

Cahiers Thomason. Bibliothèque MIR

Dually, we have strongly right enriched.

Lemma Let $\mathcal{K}, \mathcal{L}$ be two left semi-model categories, (SI, 48) and suppose $\mathcal{K} \times \mathcal{L} \xrightarrow{\otimes} \mathcal{L}$ is a strong left enrichment. p. 3.

Then:

1) If $K \in \mathcal{K}_{cof} \subseteq \mathcal{K}$, then

$K \otimes : \mathcal{L} \longrightarrow \mathcal{L}$ preserves cofibrations.

and equicofibrations = cofibrations which are weak equivalences.

2) Moreover, $K \otimes :$ preserves equivalences

(= weak equivalen

New terminology :

"weak equivalences" = equimorphisms [mixes Greek + Latin roots, but no worse than "polyvalent"]

equicofibration = cofibration + weak equivalence

equifibration = fibration and weak equivalen

Lemma Let $\mathcal{K}, \mathcal{L}$ be two left semi-model categories (SI, 48) and $\otimes : \mathcal{K} \times \mathcal{L} \longrightarrow \mathcal{L}$ a strong left enrichment. (p. 3)

Then 1) If $K \in \mathcal{K}_{cof} \subseteq \mathcal{K}$, then $K \otimes : \mathcal{L} \longrightarrow \mathcal{L}$ preserves cofibrations, equicofibrations, and those equimorphisms between cofibrant objects.

Cahiers Thomason. Bibliothèque MIR

2) If $L \in \mathcal{Z}_{cof} \subseteq \mathcal{Z}$, then $\otimes L : \mathcal{K} \to \mathcal{Z}$

preserves cofibrations, equicofibrations, and those equimorphisms between cofibrant objects.

3) The restriction of $\mathcal{K} \times \mathcal{Z} \xrightarrow{\otimes} \mathcal{Z}$ to $\mathcal{K}_{cof} \times \mathcal{Z}_{cof}$ factors thru $\mathcal{K}$ as $\mathcal{K}_{cof} \times \mathcal{Z}_{cof} \xrightarrow{\otimes_!} \mathcal{Z}_{cof} \to \mathcal{Z}$,

and $\otimes_!$ is a left enrichment in the sense of 81, 51

—

Pf:

1) As $K \in \mathcal{K}_{cof}$, $\emptyset \rightarrowtail K$ is a cofibration.

If $L \rightarrowtail L'$ is a cofibration in $\mathcal{Z}$ (resp. an equicofibration in $\mathcal{Z}$).

$$\begin{array}{ccc} K \otimes L \amalg_{\emptyset \otimes L} \emptyset \otimes L' & \longrightarrow & K \otimes L' \\ \downarrow & & \\ K \otimes L & & \end{array}$$

is a cofibration (resp. equicofibration) by axiom LSIc).

It follows from 85, 115 that $K \otimes$ preserves equimorphisms between cofibrant objects, (Brown's trick cf 80, 11).

2) similar to 1).

3) $\otimes_!$ satisfies the axiom LIa1) of 81, 51

by LSa1) and 1) $\otimes_!$ satisfies the axiom LIa2)

by LSa2) and 2)

and LSIc) implies LIc). QED.

Cahiers Thomason. Bibliothèque MIR

Lemma. Let $\mathcal{K}$, $\mathcal{L}$ be two left semi-model categories (SI, 7.8) and suppose that every object of $\mathcal{K}$ is cofibrant. Then a functor $\mathcal{K} \otimes \mathcal{L} \xrightarrow{\otimes} \mathcal{L}$ is a strong left enrichment (7.4) iff it satisfies

~~LIa1): (cf SI, 5.1) $\forall K \in \mathcal{K}$~~

sLIa1) (cf SI, 5.1)

$\forall K \in \mathcal{K} = \mathcal{K}_{cof}$, $K \otimes : \mathcal{L} \longrightarrow \mathcal{L}$ preserves initial objects, cofibrations and pushouts along a cofibration.

sLIa2) $\forall L \in \mathcal{L}$, $\otimes L : \mathcal{K} \longrightarrow \mathcal{L}$ sends $\emptyset$ to $\emptyset$, and sends cobase change along a cofibration in $\mathcal{K}$ to a cobase change in $\mathcal{L}$ (but need not preserve the cofibration).

sLIc) If $K \rightarrowtail K'$ is a cofibration in $\mathcal{K}$ and $L \rightarrowtail L'$ is a cofibration in $\mathcal{L}$, then the induced map

$$K' \otimes L \underset{K \otimes L}{\cup} K \otimes L' \longrightarrow K' \otimes L'$$

is a cofibration. It is an equicofibration if in either of the cofibrations $L \rightarrowtail L'$ or $K \rightarrowtail K'$ is also an equimorphism.

(Note the pushout is representable by axiom LB4 of SI, 7.8 as $K \otimes L \rightarrowtail K \otimes L'$ by sLIa1).)

Pf: If $\otimes$ is a strong left enrichment, the sLIa1) holds by SLIa1) and Lemma 1) 7.4. SLIa2) gives SLIa2), and SLIc) gives sLIc).

Cahiers Thomason. Bibliothèque MIR

Conversely, if SLIa1), SLIa2) or ~~SLIa1~~ SLIc) hold,

Then in the situation of $K \rightarrowtail K'$, $L \rightarrowtail L'$, SLIa1) and left model axiom LBY shows $K' \otimes L \amalg_{K \otimes L} K \otimes L'$ exists a $K \otimes L \rightarrowtail K \otimes L'$. Then SLIc) gives the rest of SLIc). SLIa2) = SLIa2), and SLIa1) is stronger than SLIa1), QED.

Danger. Watch out for colimits in $\mathcal{C}$ of us in $\mathcal{C}$.

Let R be a PID, and $\mathcal{P}$ the category of finitely generated free R-modules.

Then $\mathcal{P}$ has all finite colimits! For clearly, it has finite coproducts. It suffices to show it has coequalizers. As $\mathcal{P}$ is additive, the coequalizer of $P \underset{g}{\overset{f}{\rightrightarrows}} Q$ is the coequalizer of $P \underset{0}{\overset{f-g}{\rightrightarrows}} Q$. So it suffices to show $\mathcal{P}$ has cokernels.

Given $P \xrightarrow{f} Q$ in $\mathcal{P}$, let Q/fP be the cokernel in R-Mod. ~~A~~ ~~B~~ It is finitely generated over R. By the structure theorem for modules over a PID the quotient of Q/fP by its torsion submodule $T(Q/fP) \leq Q/fP$ is free and finitely generated.

If F is any free module and $P \xrightarrow{f} Q \xrightarrow{g} F$ with $gf = 0$, then we have an induced map $Q/fP \longrightarrow F$.

But as F is torsion free, this map factors uniquely

Cahiers Thomason. Bibliothèque MIR

$Q/fP \twoheadrightarrow (Q/fP)/T(Q/fP) \longrightarrow F.$

So the free $(Q/fP)/T(Q/fP)$ is in fact the cokernel of $P \longrightarrow Q$ in $\mathcal{P}$!!

Now let $\mathcal{C}$ be the category of bounded complexes of R-modules of finite type. Give it the left module structure where cofibrations are monomorphisms with degreewise projective cokernel (so degreewise split mono) and where equivalences are homology isomorphisms

Then $\mathcal{C}_{cof} = \mathcal{C}^b(\mathcal{P})$ $\mathcal{P}$ = f.g. projective R-modules.

$\mathcal{C} = \mathcal{C}^b$ (fg. R-modules).

So the above shows $\mathcal{C}_{cof}$ has <u>all</u> <u>finite</u> colimits, (since $\mathcal{P}$ does), but the inclusion $\mathcal{C}_{cof} \longrightarrow \mathcal{C}$ does not preserve them.

Ex.



has pushout $= 0$ in $\mathcal{C}_{cof}(\mathbb{Z})$, but pushout in

$\mathcal{C}(\mathbb{Z})$ is



Cahiers Thomason. Bibliothèque MIR

Best to hope for is $\mathcal{C}_{cof} \longrightarrow \mathcal{C}$

preserves cobase change along a cof. sn.

Works in above example.

If doesn't work in general, I'll have to be careful to always specify pushouts are to be taken in $\mathcal{C}$ in using old axiom for left enrichment (81, 51)

$$\begin{array}{ccc} C & \rightarrowtail & C' \\ \downarrow & & \\ B & & \end{array} \quad \text{in } \mathcal{C}_{cof}.$$

Take PO in $\mathcal{C}$

$$\begin{array}{ccc} C & \rightarrowtail & C' \\ \downarrow & \square & \downarrow \\ B & \longrightarrow & P \end{array}$$

the LBT $\Rightarrow \quad B \rightarrowtail P. \quad$ So $\quad \emptyset \rightarrowtail B \rightarrowtail P \quad P \in \mathcal{C}_{cf}.$

Since $\mathcal{C}_{cof}$ is a full subcat of $\mathcal{C}$, it forms P in also a pushout in $\mathcal{C}_{cof}$, hence the pushout there.

So is OK.

Lemma Let $\mathcal{C}$ be a left semi-model category (81, 48) and $\mathcal{C}_{cof}$ the full subcategory of cof. brant objects.

Then pushouts along a cof. sn are preserved by the inclusion of $\mathcal{C}_{cof} \longrightarrow \mathcal{C}$. (General pushouts need not be preserved. see example, 7-8).

Thus the inclusion $\mathcal{C}_{cof} \longrightarrow \mathcal{C}$ is a left semi-model functor (85, 29).

Cahiers Thomason. Bibliothèque MIR

Pf: Let 

be a diagram in $\mathcal{C}_{cof}$. By, the axiom LB4 (§1, 48) the pushout in $\mathcal{C}$ is $C'' \rightarrowtail C' \underset{C}{\sqcup} C''$ a cofibration.

But then $\emptyset \rightarrowtail C'' \rightarrowtail C' \underset{C}{\sqcup} C''$ gives $C' \underset{C}{\sqcup} C'' \in \mathcal{C}_{cof}$.

Since the pushout in $\mathcal{C}$ in fact lies in $\mathcal{C}_{cof}$, and $\mathcal{C}_{cof}$ is a ~~part~~ full subcategory of $\mathcal{C}$, it will also be a pushout in $\mathcal{C}_{cof}$. Since pushouts are unique up to iso, this shows $\mathcal{C}_{cof} \rightarrow \mathcal{C}$ preserves pushouts along a cofibration.

The statement that $\mathcal{C}_{cof} \rightarrow \mathcal{C}$ is a left semi-model functor now follows from the definition (§5, 29) and the method of inducing a left semi model structure on $\mathcal{C}_{cof}$ (§1, 50)

QED.

Cahiers Thomason. Bibliothèque MIR

Hypoth: Let $\mathcal{C}$ be a left semi-model category, (§1, p 48)

$\mathcal{I}_f$ the category of finite simplicial sets,

and $\mathcal{I}_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$ a strong left enrichment. (p 2).

Note as every object of $\mathcal{I}_f$ is cofibrant, we may use the equivalent formulation of the conditions on $\otimes$ given by, Lemme p. 6.

—

Def: For $\mathcal{C}$ a left semi-model category, (§1, p 48) strongly left enriched over $\mathcal{I}_f$ (p 2), define

cf §5 p 110.

$$\mathrm{Hom} : \mathcal{C}^{op} \times \mathcal{C} \longrightarrow \Delta^{op}\text{-sets}$$

by $\mathrm{Hom}(C, C')_p = \mathrm{Mor}_{\mathcal{C}}(\Delta[p] \otimes C, C')$

—

Ask if $\mathrm{Mor}_{\Delta^{op}\text{Sets}}(K, \mathrm{Hom}(C, C')) \cong \mathrm{Mor}_{\mathcal{C}}(K \otimes C, C')$

—

ok : $K = \Delta\Sigma[1]$.

in general need $K \cong \varinjlim_{n\text{d}\,\Delta/K} \Delta \quad \Rightarrow \quad K \otimes C \cong \varinjlim_{n\text{d}\,\Delta/K} (\Delta[?] \otimes C)$.

ie we

$\otimes C : \mathcal{I}_f \longrightarrow \mathcal{C}$ preserves finite colimits.

—

If C cofibrant ok by §2 p 5 (cf §5 p 110)
§2, 6B

Cahiers Thomason. Bibliothèque MIR

It looks like this argument can in fact be generalized to work with $\mathcal{C}$ not cof. stat.

Key point is for a general $\mathcal{C}$, given

$$\begin{array}{ccc} K & \longrightarrow & L \\ \downarrow & \text{po} & \downarrow \\ N & \longrightarrow & P \end{array} \quad \text{in } \mathcal{A}_f$$

$$\begin{array}{ccc} K \otimes C & \longrightarrow & L \otimes C \\ \downarrow & & \downarrow \\ N \otimes C & \longrightarrow & P \otimes C \end{array}$$

is a pushout square in $\mathcal{C}$, ~~etc.~~ (axiom LS[a1]),

even though $K \otimes C \to L \otimes C$ need not be a cofibration if $\mathcal{C}$ not cofstat.

In ~~proof~~ particular $\forall N$ ~~$\coprod_N \Delta[0] \to \coprod_N \Delta[0]$~~

$\coprod_N (\Delta[0] \otimes C)$ exists in $\mathcal{C}$ as a ~~sum~~ coproduct

$\left(= \left(\coprod_N \Delta[0] \right) \otimes C \right)$.

—

But this is getting complicated.

By the time we add the extra hypotheses of ~~p.37~~ 136 we'll wind up with an enrichment over Δ° sets ~~anyway~~ anyway.

Cahiers Thomason. Bibliothèque MIR

So just go for it. Will be simpler.

—

~~Natural~~ Hypotheses:

$(\mathcal{C}, \text{cof.s}\,\mathcal{C}, \text{equi}\,\mathcal{C}, \text{f.s}\,\mathcal{C})$ is a basic model category with functorial factorization

~~Natural~~ Hypotheses ~~Enrich~~ Rich Hypotheses

$(\mathcal{C}, \text{cof.s}\,\mathcal{C}, \text{equi}\,\mathcal{C}, \text{f.s}\,\mathcal{C})$ a basic model category

i.e. satisfies axioms B1 – B7 of 78, 156-157.

Surrose $\mathcal{C}$ is enriched over Δ^{op}-sets in the usual sense of category theory:

$$\underline{\text{Hom}} : \mathcal{C}^{op} \times \mathcal{C} \longrightarrow \Delta^{op}\text{-sets} \quad \text{s.t. } \dots$$

Let $\mathcal{I}_f \subseteq \Delta^{op}$-sets be the full subcategory of those simplicial sets with only finitely many non-degenerate simplices (i.e. "finite simplicial sets")

We surrose further that there are functors — Gepka notes, p. 19

$$\mathcal{I}_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C} \qquad \mathcal{I}_f^{op} \times \mathcal{C} \xrightarrow{\pitchfork} \mathcal{C}$$

~~Map(,)~~ ~~$\mathcal{I}_f^{op}$~~

such that there are natural isomorphisms for $K \in \mathcal{I}_f$, $C, C' \in \mathcal{C}$

$$\underline{\text{Hom}}(K \otimes C, C') \cong \text{Map}(K, \underline{\text{Hom}}(C, C')) \cong \underline{\text{Hom}}(C, K \pitchfork C')$$

Cahiers Thomason. Bibliothèque MIR

Moreover, we suppose

~~LE~~ ~~If $K \rightarrowtail K'$ is a cofibration in $\mathcal{I}_f$ and $C \rightarrowtail C'$ is a cofibration in $\mathcal{C}$~~



LE1 $\forall K$, $\mathcal{I}_f$, $K \otimes : \mathcal{C} \longrightarrow \mathcal{C}$ preserves cofibrations.

LE2 If $K \rightarrowtail K'$ is a cofibration in $\mathcal{I}_f$ and $C \rightarrowtail C'$ is a cofibration in $\mathcal{C}$, then the induced map $K' \otimes C \underset{K \otimes C}{\sqcup} K \otimes C' \longrightarrow K' \otimes C'$ is a cofibration in $\mathcal{C}$.

LE If $K \rightarrowtail K'$ is a cofibration in $\mathcal{I}_f$ and $C \rightarrowtail C'$ is a cofibration in $\mathcal{C}$, then the pushout $K' \otimes C \underset{K \otimes C}{\sqcup} K \otimes C'$ is representable in $\mathcal{C}$, and the induced map $K' \otimes C \underset{K \otimes C}{\sqcup} K \otimes C' \longrightarrow K' \otimes C'$ is a cofibration in $\mathcal{C}$.

It is a equivalence if either of the cofibrations is an equivalence.

Remark: The hypothesis that $K' \otimes C \underset{K \otimes C}{\sqcup} K \otimes C'$ is representable is in fact redundant. For $\emptyset \otimes C \cong \emptyset$ (follows from adjunct. $\underline{\mathrm{Hom}}\, \mathrm{Mor}_{\mathcal{C}}(\emptyset \otimes C, C') = \underline{\mathrm{Hom}}(\emptyset \otimes C, C')_0 = \mathrm{Map}(\emptyset, \underline{\mathrm{Hom}}(C, C'))_0 = *$.)

and $\forall K'$ $K' \otimes C \underset{\emptyset}{\sqcup} \emptyset = K' \otimes C$. ~~Thus we set LE =>~~

Since $\forall K'$, $\emptyset \rightarrowtail K' \in \mathcal{I}_f$, LE gives this instance of LE gives $K' \otimes$ preserves cofibrations.

Cahiers Thomason. Bibliothèque MIR

But then in the general case $C \rightarrowtail C' \Rightarrow K'\otimes C \rightarrowtail K'\otimes C'$,

and now axiom B4 for basic model cats (or LB4 for left model cat)

gives the pushout exist in general.)

RE If $K \rightarrowtail K'$ is a cofibration in Δ_f and $C \twoheadrightarrow C'$ is a fibration in $\mathcal{C}$, then

$$K' \overset{\Delta}{\pitchfork} C' \underset{K \overset{\Delta}{\pitchfork} C'}{\times} K \overset{\Delta}{\pitchfork} C \quad \text{is represented in } \mathcal{C}$$

and the induced map

$$K' \overset{\Delta}{\pitchfork} C \longrightarrow K' \overset{\Delta}{\pitchfork} C' \underset{K \overset{\Delta}{\pitchfork} C'}{\times} K \overset{\Delta}{\pitchfork} C \quad \text{is a fibration.}$$

This map is an equifibration if either the cofibration $K \rightarrowtail K'$ or the fibration $C \twoheadrightarrow C'$ is an equimorphism.

[the dual to remark LE applies to RE.]

Lemma: If $(\mathcal{C}, \mathrm{cof}\,\mathcal{C}, \mathrm{equi}\,\mathcal{C}, \mathrm{fib}\,\mathcal{C}, \underline{\mathrm{Hom}}, \otimes, \overset{\Delta}{\pitchfork})$ satisfies the hypotheses of p 13, then the dual model category $(\mathcal{C}^{op}, \mathrm{cof}\,\mathcal{C}^{op} = (\mathrm{fib}\,\mathcal{C})^{op}, \mathrm{equi}\,\mathcal{C}^{op} = (\mathrm{equi}\,\mathcal{C})^{op}, \mathrm{fib}\,\mathcal{C}^{op} = (\mathrm{cof}\,\mathcal{C})^{op})$ (cf 78, p 159)

with $\underline{\mathrm{Hom}}^{op} : (\mathcal{C}^{op})^{op} \times \mathcal{C}^{op} = \mathcal{C} \times \mathcal{C}^{op} \cong \mathcal{C}^{op} \times \mathcal{C} \xrightarrow{\underline{\mathrm{Hom}}} \Delta^{op}$-sets

(i.e. $\underline{\mathrm{Hom}}^{op}(C', C) = \underline{\mathrm{Hom}}(C, C')$, etc.).

[cf Eilenberg–Kelley III 2.1]

~~$K\otimes\mathcal{C}^{op}$~~ $\Delta_f \times \mathcal{C}^{op} \xrightarrow{\otimes^{op}} \mathcal{C}^{op}$

$$= (\Delta_f^{op} \times \mathcal{C})^{op} \xrightarrow{\overset{\Delta}{\pitchfork}} \mathcal{C}^{op},$$

$$\overset{\Delta}{\pitchfork}{}^{op} : \Delta_f^{op} \times \mathcal{C}^{op} \longrightarrow \mathcal{C}^{op}$$
$$\| \qquad\qquad \|$$
$$(\Delta_f \times \mathcal{C})^{op} \xrightarrow{\otimes} \mathcal{C}^{op}$$

Cahiers Thomason. Bibliothèque MIR

satisfies the ↑rich hypotheses of 1.3.

Thus the ↑rich hypotheses are invariant under duality.

—

Proof Suppose the rich hypotheses of 1.3, except LE on RE.

Then

1) There is a natural identification preserving composition and unit

~~$[\underline{\mathrm{Hom}}(C, C')]_0 = \mathrm{Map}(E^{op} \times E, Kan)$~~

~~Hom~~

$$[\underline{\mathrm{Hom}}(C, C')]_0 \cong \mathrm{Map}_e(C, C')$$

~~2) There are compatible natural isomorphisms compa~~

2) There are natural isomorphisms

$$\Delta[0] \otimes C \cong C \qquad \Delta[0] \pitchfork C' \cong C'$$

render commutative the diagram

$$\left[\underline{\mathrm{Hom}}(\Delta[0] \otimes C, C')\right]_0 \cong \left[\mathrm{Map}(\Delta[0], \underline{\mathrm{Hom}}(C, C'))\right]_0 \cong \left[\underline{\mathrm{Hom}}(C, \Delta[0] \pitchfork C')\right]_0$$

$$\| \qquad\qquad \| \qquad\qquad \|$$

$$\mathrm{Map}_e(\Delta[0] \otimes C, C') \qquad \left[\underline{\mathrm{Hom}}(C, C')\right]_0 \qquad \mathrm{Map}_e(C, \Delta[0] \pitchfork C')$$

$$\cong \qquad\qquad \| \qquad\qquad \cong$$

$$\mathrm{Map}_e(C, C')$$

Cahiers Thomason. Bibliothèque MIR

3) If $K : \underline{D} \longrightarrow \mathcal{I}_f$ is any diagram such that the colimit taken in Δ^{op}-sets $\varinjlim_{\underline{D}} K$ lies in fact in $\mathcal{I}_f$, then for all $C \in \mathcal{C}$ ~~the natural~~ ~~map~~ than $(\varinjlim_{\underline{D}} K) \otimes C$ is the colimit in $\mathcal{C}$ of the diagram $K \otimes C : \underline{D} \longrightarrow \mathcal{C}$.

(Maybe $\mathcal{I}_f \longrightarrow \Delta^{op}$-sets preserves all colimits that exist in $\mathcal{I}_f$??).

Yes but proof ugly. Not worth it.

Dually if $L : \underline{D}^{op} \longrightarrow \mathcal{I}_f$ is any diagram such that the colimit in Δ^{op}-set $\varinjlim_{\underline{D}^{op}} L$ lies in fact in $\mathcal{I}_f$, then $\forall C' \in \mathcal{C}$ ~~$\varinjlim_{\underline{D}^{op}}$~~ $(\varinjlim_{\underline{D}^{op}} L) \pitchfork C'$ represents the limit of $\underline{D} \longrightarrow \mathcal{C}$. $L \pitchfork C' : \underline{D} \longrightarrow \mathcal{C}$.

4) If ~~$K : \mathcal{I}_f$~~ $F : \underline{D} \longrightarrow \mathcal{C}$ a is any diagram with a colimit $\varinjlim_{\underline{D}} F$ in $\mathcal{C}$, then $\forall K \in \mathcal{I}_f$ ~~the natural map~~ $K \otimes (\varinjlim_{\underline{D}} F)$ represents the colimit of $K \otimes F : \underline{D} \longrightarrow \mathcal{C}$ $(D \longmapsto K \otimes FD)$.

Dually, if $F' : \underline{D} \longrightarrow \mathcal{C}$ is any diagram with a limit $\varprojlim_{\underline{D}} F'$ in $\mathcal{C}$, then $\forall K \in \mathcal{I}_f$ $K \pitchfork (\varprojlim_{\underline{D}} F')$ represents the limit of $K \pitchfork F' : \underline{D} \longrightarrow \mathcal{C}$.

Cahiers Thomason. Bibliothèque MIR

Pf 1) Follows from

$$[\underline{\mathrm{Hom}}(C,C')]_0 = \mathrm{Map}(X,Y) \qquad \mathrm{Mor}_{\Delta^{op}\mathrm{Set}}(X,Y)$$

and standard properties of enrichment over a ~~cat~~ symmetric ~~closed category~~ monoidal closed category. (cf Kelly, Basic ~~Concepts~~ Concepts of Enriched Category Theory 1.3)

2) We have obtained from the adjunction isos natural isos

$$\mathrm{Mor}_{\mathcal{C}}(C \otimes \Delta[0], C') \cong \mathrm{Mor}_{\mathcal{C}}(C, C') \cong \mathrm{Mor}_{\mathcal{C}}(C, \Delta[0] \pitchfork C').$$

That these are induced by unique ~~isos~~ natural isos

$$C \otimes \Delta[0] \cong C, \qquad C' \cong \Delta[0] \pitchfork C'$$

now follows from Yoneda

3) ~~$\underline{\mathrm{Hom}}(K \otimes C, C') \cong \mathrm{Mor}(K, \underline{\mathrm{Hom}}(C, C'))$~~

$$\underline{\mathrm{Hom}}(K \otimes C, C') \cong \mathrm{Map}(K, \underline{\mathrm{Hom}}(C, C'))$$

implies on taking Γ_0 and using 1)

a natural iso

$$\mathrm{Mor}_{\mathcal{C}}(K \otimes C, C') \cong \mathrm{Mor}_{\Delta^{op}\mathrm{Set}}(K, \underline{\mathrm{Hom}}(C, C')).$$

Then $\mathrm{Mor}_{\mathcal{C}}((\varinjlim_{\mathcal{D}} K) \otimes C, C') \cong \mathrm{Mor}_{\Delta^{op}\mathrm{Set}}(\varinjlim_{\mathcal{D}} K, \underline{\mathrm{Hom}}(C, C'))$

$$\varprojlim_{\mathcal{D}^{op}} \mathrm{Mor}_{\mathcal{C}}(K \otimes C, C') \cong \varprojlim_{\mathcal{D}^{op}} \mathrm{Mor}_{\Delta^{op}\mathrm{Set}}(K, \underline{\mathrm{Hom}}(C, C'))$$

Cahiers Thomason. Bibliothèque MIR

This shows $(\varinjlim_{\mathcal{D}} K) \otimes C$ represents the colimit, by definition.

(Note $K \otimes C$ as $I_f \longrightarrow C$ is not left adjoint to $\underline{Hom}(C, \ )$, as the latter goes to Δ^{op} Sets, not to I_f !)

Dual half of 3) is similar, & or follows by duality!

4) We have

$$Mor_{\mathcal{C}}(K \otimes C, C') \cong Mor_{\Delta^{op}\text{s}}(K, \underline{Hom}(C, C')) \cong Mor_{\mathcal{C}}(C, K \pitchfork C')$$

So $K \otimes$ is left adjoint to $K \pitchfork : \mathcal{C} \longrightarrow \mathcal{C}$, and hence preserves colimits. (or argue as in part 3)).

Dually, $K \pitchfork$ preserves whatever limits that may exist

QED

Notation. $K \pitchfork X$ looks too symmetric, and is clunky.

$Mor(K, X)$ too long-winded, and obscures duality.

Better candidates (exist in TeX

(Symbols which exist in AmsTeX,

$\supset$	\Supset	all binary relations
$\sqsupset$	\sqsupset	(available binary operators too symmetric)
$\triangleright$	\vartriangleright	
$\propto$	\varpropto	
$\blacktriangleright$	\blacktriangleright	all in Further Funb. use \Use AMS symbols

$\propto$	\propto	standard ams fm
$\nsubseteq$		
$\sqsubseteq$	\sqsubseteq	

Cahiers Thomason. Bibliothèque MIR

$\sqsupset$ \sqsupset available in latexsym package.

$\sqsupseteq$ \sqsupseteq in plain latex.

$\sqsupseteq$ \sqsupseteq - TeX.

~~TeX has \triangleright a~~

TeX (

$\triangleright$ = \triangleright ~~also~~, even in binary op.

I. ~~Note~~ Latex

Binop $\rhd$ = \rhd needs latexsym package

~~Amsfonts~~.

Amsfonts seems not to have a bin $\rhd$ as binary operator

But could use $\vartriangleright$ \vartriangleright or

Conclusion

Write ~~adj~~ enriched adjoint.

$$Map(K, \underline{Hom}(C, C')) \cong \underline{Hom}(C, K \triangleright C')$$

~~where~~ $K \triangleright C'$ is the cotensor $(K_f)^{ob} \times C \xrightarrow{\triangleright} C$

$A \triangleright B$ i.e. $Map(A, B) = \{A \longrightarrow B\}$.

In Latex ~~to~~ use binary operator $\rhd$ = \rhd
in ~~the~~ latexsym package.

OK for AMS Latex ?

In ~~TeX~~ AMS TeX, use binary relation $\triangleright$ = \vartriangleright
~~defined~~ retyped to be a binary operator.

Cahiers Thomason. Bibliothèque MIR

In plain Tex use the dinky binary operat. $\triangleright$ = \triangleright.

Arghh But maybe $\Delta[n] \triangleright X$ will look bad.

If so fall back to $\Delta[n] \pitchfork X$

Lemma: Let $(\mathcal{C}, \text{cofib}\,\mathcal{C}, \text{equiv}\,\mathcal{C}, \text{fib}\,\mathcal{C}, \underline{\text{Hom}}, \otimes, \triangleright)$ satisfy the ~~hypotheses~~ rich hypotheses of §3 except for possibly LE and RE.

~~Then LE and RE~~

Then the ~~conclusions~~ LE and RE are each seperately equivalent to

EB: If $C \rightarrowtail C'$ is a cofibration in $\mathcal{C}$ and $D \twoheadrightarrow D'$ is a fibration the induced map

$$\underline{\text{Hom}}(C', D) \longrightarrow \underline{\text{Hom}}(C, D) \times_{\underline{\text{Hom}}(C, D')} \underline{\text{Hom}}(C', D')$$

EB: If $C \rightarrowtail C'$ is a cofibration in $\mathcal{C}$ and $D' \twoheadrightarrow D$ is a fibration in $\mathcal{C}$, the induced map

$$\underline{\text{Hom}}(C', D') \longrightarrow \underline{\text{Hom}}(C, D') \times_{\underline{\text{Hom}}(C, D)} \underline{\text{Hom}}(C', D)$$

is a Kan fibration in Δ^{op}sets.

It is a "weak equivalence" if either the cofibration $C \rightarrowtail C'$ or the fibration $D' \twoheadrightarrow D$ is also an equivalence.

Pf: In particular LE holds iff RE holds.

Cahiers Thomason. Bibliothèque MIR

No Quillen's version (Homotopical Algebra II 2.3 Prop 3)

depends on characterizing fibrations as morphisms with the right lifting property for equicofibrations etc

—

What I have is

cf Quillen Homotopical Algebra

Lemma: Let $(\mathcal{C}, \mathrm{cof}\mathcal{C}, \mathrm{equi}\mathcal{C}, \mathrm{fib}\mathcal{C})$ ~~basic model~~ category satisfy B1 – B6 of the basic model axioms (ie no factorization axiom) § 78, 156

Let $\mathcal{C}$ be enriched over Δ°-sets via $\underline{\mathrm{Hom}}$, and suppose $\mathcal{C}$ has a tensor $\otimes$ and cotensor $\triangleright$ over $I_f \subseteq \Delta^{\circ}$-Sets s.t. there are natural isomorphisms

for $K \subseteq I_f$, $C, C' \in \mathcal{C}$

$$\underline{\mathrm{Hom}}(K \otimes C, C') \cong \mathrm{Map}(K, \underline{\mathrm{Hom}}(C, C')) \cong \underline{\mathrm{Hom}}(C, K \triangleright C)$$

Then the ~~combination~~ conjunction of axioms LE and RE (p. 14) is equivalent to the following:

EB If $C \rightarrowtail C'$ is a cofibration in $\mathcal{C}$ and $D' \twoheadrightarrow D$ is a fibration, then the induced map

$$\underline{\mathrm{Hom}}(C', D') \longrightarrow \underline{\mathrm{Hom}}(C, D') \times_{\underline{\mathrm{Hom}}(C, D)} \underline{\mathrm{Hom}}(C', D)$$

is a Kan fibration of simplicial sets. Moreover if "a "weak equivalence" in the usual sense if either the cofibration $C \rightarrowtail C'$ or the fibration $D' \twoheadrightarrow D$ is an equimorphism in $\mathcal{C}$.

Cahiers Thomason. Bibliothèque MIR

Pf:

First note $f : A \to B$ in Δ^{op}-sets, the adjunction implies a bijective correspondence between solid diagrams (solid or solid plus dotted arrows)

$$\begin{array}{ccc} A & \longrightarrow & \underline{Hom}(C', D') \\ \downarrow & \nearrow & \downarrow \\ B & \longrightarrow & \underline{Hom}(C, D') \times \underline{Hom}(C', D) \\ & & \underline{Hom}(C, D) \end{array}$$

and



ie



both the adjunction!

Similarly, these correspond injectively to



Cahiers Thomason. Bibliothèque MIR

cf §5 p.112
Quillen Homotopical Alg. Prop 3
II 2.3

Lemma Let $(\mathcal{C}, \mathrm{cof}\mathcal{C}, \mathrm{equi}\mathcal{C}, \mathrm{fib}\mathcal{C})$ satisfy B1 – B6 of §7 p.156, ie the basic model axioms except factorization.

Suppose $\mathcal{C}$ is enriched over Δ^{op}-sets, with internal Hom. $\mathcal{C}^{op} \times \mathcal{C} \longrightarrow \Delta^{op}$-sets.

Suppose there is a functor

$$\mathcal{A}_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$$

and a natural isomorphism

for $K \in \mathcal{A}_f$, $C, C' \in \mathcal{C}$

$$\underline{\mathrm{Hom}}(K \otimes C, C') \cong \underline{\mathrm{Map}}(K, \underline{\mathrm{Hom}}(C, C'))$$

If $\otimes$ satisfies LE (p.141), then

BE:

for each cofibration $C \rightarrowtail C'$ in $\mathcal{C}$ and each fibration $D' \twoheadrightarrow D$ in $\mathcal{C}$, the induced map

$$\underline{\mathrm{Hom}}(C', D') \longrightarrow \underline{\mathrm{Hom}}(C, D') \times_{\underline{\mathrm{Hom}}(C, D)} \underline{\mathrm{Hom}}(C', D)$$

is a Kan fibration of simplicial sets.

It is a weak equivalence in the usual sense if either the cofibration $C \rightarrowtail C'$ or the fibration $D' \twoheadrightarrow D$ is an equimorphism.

(Dually, if there is a $\mathcal{A}_f^{op} \times \mathcal{C} \xrightarrow{\triangleright} \mathcal{C}$ such that $\underline{\mathrm{Map}}(K, \underline{\mathrm{Hom}}(C, C')) \cong \underline{\mathrm{Hom}}(C, K \triangleright C')$ naturally, and $\triangleright$ satisfies RE, then EB holds.

Cahiers Thomason. Bibliothèque MIR

Pf: The dual statement indeed is dual, ie the given statement for $\mathcal{C}^{op}$.

So it suffices to prove the given statement.

The ~~adjunction~~ isomorphism

$$\underline{\mathrm{Hom}}(K \otimes C, C') \cong \underline{\mathrm{Hom}}(K, \underline{\mathrm{Hom}}(C, C'))$$

implies on taking $[\ ]_0$ on both

$$\mathrm{Hom}_{\mathcal{C}}(K \otimes C, C') \cong \mathrm{Hom}_{\Delta^{op}\mathrm{Set}}(K, \underline{\mathrm{Hom}}(C, C')).$$

This induces a natural bijection between diagrams of solid arrows (resp. solid and dotted arrows) in ~~$\mathcal{C}^{op}$~~ ~~from $K \to \underline{\mathrm{Hom}}(C, D)$~~

$K \to L$ in Δ^{op}-set.



cf §5, 118



ie



Cahiers Thomason. Bibliothèque MIR

Now to say $\underline{Hom}(C', D') \longrightarrow \underline{Hom}(C, D') \times_{\underline{Hom}(C,D)} \underline{Hom}(C', D)$

is a Kan fibration is to say for all horn $\Lambda[n] \xrightarrow{\sim} \Delta[n]$

one can always find a ~~sol~~ dotted arrow filling in an

solid arrow diagram of the form

$$\begin{array}{ccc} \Lambda[n] & \longrightarrow & \underline{Hom}(C', D') \\ \sim\downarrow & \nearrow & \downarrow \\ \Delta[n] & \longrightarrow & \underline{Hom}(C, D') \times_{\underline{Hom}(C,D)} \underline{Hom}(C', D) \end{array}$$

But the corresponding diagram is

$$\begin{array}{ccc} \Delta[n] \otimes C \sqcup_{\Lambda[n]\otimes C} \Lambda[n] \otimes C' & \longrightarrow & D' \\ \downarrow & \nearrow & \downarrow \\ \Delta[n] \otimes C' & \longrightarrow & D \end{array}$$

By LE, the left vertical map is a equivalence.

Then by axiom B6, the fill-in arrow exists. Thus

the fill-in exists in the corresponding diagram.

~~If~~ Suppose now either $C \hookrightarrow C'$ or $D' \twoheadrightarrow D$ is an

equivalence. To show the ~~Kan fibration~~ map is Δ^n-sat

$$\underline{Hom}(C', D') \longrightarrow \underline{Hom}(C, D') \times_{\underline{Hom}(C,D)} \underline{Hom}(C', D)$$

is a Kan fibration and weak equivalence

Cahiers Thomason. Bibliothèque MIR

it suffices to show ~~an solid arrow diagram of $\dot\Delta[n] \to \Delta[n]$~~ for all n on $\dot\Delta[n] \to \Delta[n]$ that any solid arrow diagram of the form

$$\begin{array}{ccc} \dot\Delta[n] & \longrightarrow & \underline{\mathrm{Hom}}(C', D') \\ \downarrow & \nearrow & \downarrow \\ \Delta[n] & \longrightarrow & \underline{\mathrm{Hom}}(C, D') \times_{\underline{\mathrm{Hom}}(C, D)} \underline{\mathrm{Hom}}(C', D) \end{array}$$

admits a dotted fill-in. (cf Quillen Homotopical Algebra II 2.2 Def 1, II 3.4 Thm 3, II 3.19 Prop 7)

But the corresponding diagram

$$\begin{array}{ccc} \Delta[n] \otimes C \cup_{\dot\Delta[n]\otimes C} \dot\Delta[n] \otimes C' & \longrightarrow & D' \\ \downarrow & \nearrow & \downarrow \\ \Delta[n] \otimes C' & \longrightarrow & D \end{array}$$

~~But~~ LE gives that $\Delta[n] \otimes C \cup_{\dot\Delta[n]\otimes C} \dot\Delta[n] \otimes C' \longrightarrow \Delta[n] \otimes C'$ is a cofibration, and an equicofibration ~~of~~ if $C \rightarrowtail C'$ is an equimorphism. Thus either the left cofibration or the right fibration $D' \twoheadrightarrow D$ is an equimorphism, and the dotted arrow exists by axiom BM6.

QED.

Remark: Conversely, BE implies $\forall$ $K \rightarrowtail L$ "Δ^{op}-set" ~~that~~ $\forall$ $C \rightarrowtail C'$ in $\mathcal{C}$, the ~~map~~ ~~is~~ ~~a~~ ~~fib~~ ~~if~~

Cahiers Thomason. Bibliothèque MIR

~~K~~ "$L \otimes C \underset{K \otimes C}{\sqcup} K \otimes C'$" is representable in $\mathcal{C}$, then

$L \otimes C \underset{K \otimes C}{\sqcup} K \otimes C' \longrightarrow L \otimes C'$ has the left

lifting property for trivial fibrations.

But with my basic model axioms, ~~th~~ unlike Quillen's,

this does not imply $L \otimes C \underset{K \otimes C}{\sqcup} K \otimes C' \to L \otimes C'$ is a

cofibration.

So I can't deduce LE from EB.

(cf Quillen, Homotopical Algebra II 2.3 Prop 3)

—

cf. 38

cf 54(a) one-sided

Lemma:

Let $(\mathcal{C}, \text{cof}\,\mathcal{C}, \text{equ}\,\mathcal{C}, \text{fib}\,\mathcal{C})$ be a basic model category

with functorial factorization

(ie satisfies B1–B6 of a F1, F1' of 78 . 156–158 ; here B7

note this is slight reduction in presence of enrichment below, which should

allow construction of cylinders, cf. cf 79, 72 C1 + enrich ⇒ C2 ⇒ F1, F1' ?)

Suppose $\mathcal{C}$ is enriched over Δ^{op}-set by Hom,

and $\exists \quad \Delta_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$ or $\Delta_f^{op} \times \mathcal{C} \xrightarrow{\triangleright} \mathcal{C}$

with natural isomorphisms

$$\underline{\text{Hom}}(K \otimes C, C') \cong \text{Mor}(K, \underline{\text{Hom}}(C, C')) \cong \underline{\text{Hom}}(C, \mathcal{C}' K \triangleright C')$$

~~with the~~

and s.t. RE or LE cf. p 14–15 hold

(ie $\mathcal{C}$ satisfies the Rich Hypotheses of p 13)

Cahiers Thomason. Bibliothèque MIR

~~Suppose that $\mathcal{C}$ is~~

Let λ be an infinite cardinal, and suppose that $\mathcal{C}$ is λ-productive (resp. strongly λ-productive) and ω-summable (resp. strongly ω-summable).

(SI, p. 118, p. 117).

Then

The underlying right model category of $\mathcal{C}$ w.r.t. $\mathbb{1}_f^{\omega} \times \mathcal{C} \xrightarrow{\triangleright} \mathcal{C}$

satisfies

a) There is a natural isomorphism $\forall\ C$

$$C \cong \Delta[\omega] \triangleleft C$$

b) For all indexing sets I with $\#I < \lambda$, and all families $\{C_i\}_{i \in I}$ in $\mathcal{C}_{f.s.}$ (resp. $\mathcal{C}$)

and all $K \in \mathbb{1}_f$ the natural map

$$K \triangleright \left(\prod_I C_i\right) \longrightarrow \prod_I (K \triangleright C_i)$$

is an isomorphism.

c) $\forall\ C, C' \in \mathcal{C}$, $\forall\ K, L \in \mathbb{1}_f$ there are natural isomorphisms

$$K \triangleright (L \triangleright C)$$

SII

$$((K \times L) \triangleright C)$$

SII

$$(L \times K) \triangleright C$$

SII

$$L \triangleright (K \triangleright C)$$

Cahiers Thomason. Bibliothèque MIR

c) $\forall\, C \in \mathcal{C}$, $\forall K, L \in \mathcal{I}_c$ are in a natural isomorphism

$$K \triangleright (L \triangleright C) \cong (K \times L) \triangleright C.$$

This isomorphism and the symmetry of $\times$ in $\mathcal{I}_c$ induces

$$\gamma : K \triangleright (L \triangleright C) \cong L \triangleright (K \triangleright C)$$

satisfying the requirements of §5, 141 NEX, §5, 150 c).

The iso $K \triangleright (L \triangleright C) \cong (K \times L) \triangleright C$ satisfies the commutativity condition of §5, 149 (dual to extra hypotheses §5, 136).

d) If $\cdots \to C_n \to \cdots \to C_0$ is a tower of fibrations in $\mathcal{C}_{f.s}$ (resp. in $\mathcal{C}$), then $\forall K \in \mathcal{I}_c$ the natural map

$$K \triangleleft (\varprojlim_{\omega} C_i) \longrightarrow \varprojlim_{\omega} (K \triangleleft C_i)$$

is an isomorphism

e) Then $\mathcal{C}$ with $\triangleleft : \mathcal{I}_c^{op} \times \mathcal{C}_{f.s} \to \mathcal{C}_{f.s}$

satisfies: the duals of

the weak hypotheses of §1, 143

the strong hypotheses of §3, 143

and the extra hypotheses of §5, 136

Cahiers Thomason. Bibliothèque MIR

Pf: a) " Prop 2) p16.

b) ~~If all C_i are~~

For $C_i \in \mathcal{C}_0$ we have $\emptyset \to K$, $C_i \to *$

By RE this gives $K \triangleright C_i \to \emptyset \triangleright C_i \times_{\emptyset \triangleright *} K \triangleright * \simeq * \times_{*} * = *$

is a fibration, so $K \triangleright C_i$ is fibrant. Thus ~~if~~ in the case of $\mathcal{K}$-products, all the C_i, hence all $K \triangleright C_i$ are fibrant, so

~~If~~ $\prod_I K \triangleright C_i$ exists. Then the case of strong $\mathcal{K}$-products

for an $\{C_i\}_{i \in I}$ $\#I \in \mathcal{K}$, $\prod_I K \triangleright C_i$ exists.

not really need

or the implies exists.

Now Prop 4) p16 gives $K \triangleright (\prod_I C_i)$ represented $\prod_I (K \triangleright C_i)$

So the canonical map is an iso.

c) For all $C' \in \mathcal{C}$

$$\underline{\mathrm{Hom}}(C', K \triangleright (L \triangleright C)) \cong \underline{\mathrm{Map}}(K, \underline{\mathrm{Hom}}(C', L \triangleright C))$$

$$\cong \underline{\mathrm{Map}}(K, \underline{\mathrm{Map}}(L, \underline{\mathrm{Hom}}(C', C))$$

$$\cong \underline{\mathrm{Map}}(K \times L, \underline{\mathrm{Hom}}(C', C))$$

$$\cong \underline{\mathrm{Hom}}(C', (K \times L) \triangleright C)$$

Taken $\lceil \rceil_0$ we get $K \triangleright (L \triangleright C)$ or $(K \times L) \triangleright C$ representing the same functor, so they are isomorphic by ~~Yoneda~~ Yoneda.

The compatibilities required by $\underline{\S 5}$ p41 NEX, $\underline{\S 5}$ p50 c) and $\underline{\S 5}$ p49 follow from Prop 2) p16 and computations in the ~~closed~~ symmetric monoidal closed model category $\underline{\Delta^{op}\text{-sets}}$

Cahiers Thomason. Bibliothèque MIR

d) Follows from Prop 4) p 16.

e) As to the dual weak hypotheses §1 p 143

1) follows by forgetting the dual part the model structure (cf the statement of 1) §1 p 143 !!

2) ~~The enrich~~

If $K \in \mathcal{I}_f$ $C \in \mathcal{C}_{cof}$ or $\emptyset \to K$ ($\mathcal{I}_{f\,cof} = \mathcal{I}_f$)

or $\emptyset \to C$ as in LE so $\emptyset \to K \otimes C$ ~~Reject~~.

so $K \otimes C \in \mathcal{C}_{cof}$ (not $\emptyset \otimes C = \emptyset = K \otimes \emptyset$ by Prop 3), 4) p 16

~~Then the~~

so $\mathcal{I}_f \times \mathcal{C}_{cof} \xrightarrow{\otimes} \mathcal{C}_{cof}$.

Now the left enriched axioms follow (cf §1 p 51

LIa1) cf §1 by Prop 3) p 16 + LE gives $\forall K \in \mathcal{I}_f$ $\emptyset \to K$

LIa2) follows by Prop 4) p 16 + LE gives $\forall C \in \mathcal{C}_{cof}$, $\emptyset \to C$.

LIc) is just LE.

Ann C ...)(-

e) As to the duals of

the weak hypotheses of §1 p 143 (cf §5 p 145)

1) holds by just forgetting the left part of the model structure (cf the very statement of §1 p 143 #1).

2) If $K \in \mathcal{I}_f$ or $C \in \mathcal{C}_{f.s.}$, $K \in \mathcal{I}_f = \mathcal{I}_{f\,cof}$ so $\emptyset \to K$

or $C \to *$. Then RE gives $K \pitchfork C \to K \pitchfork * \times_{\emptyset \pitchfork *} \emptyset \pitchfork C$

As $K \pitchfork * = * = \emptyset \pitchfork *$? by Prop 16 3)+4)

Cahiers Thomason. Bibliothèque MIR

we have $K \triangleright C \longrightarrow +$, i.e. $K \triangleright C \in \mathcal{C}_{f.s.}$

The D restriction is

$$(1_f)^\circ \times \mathcal{C}_{f.s} \longrightarrow \mathcal{C}_{f.s.}$$

This restriction satisfies the right enrichment axioms (§1, p 51) dualized as in §5, p 146.

RIa1) holds since ~~$\Phi \in K$~~ $\forall K \in 1_f$ by Prop 16 4) and RE

RIa2) holds since $\forall C \in \mathcal{C}_{f.s.}$ $C \longrightarrow +$ by Prop 16 3) and RE.

RIb) is emph

RIc) is just RE.

3) and 4) hold since we assume $\mathcal{C}$ is ω-summable and $\aleph_0$-productive.

5. This verifies the weak hypotheses.

The strong hypotheses of §3, p 143 dualized (cf §5, p 148) just adds to the weak hypothesis

3) = RWN : $C \xrightarrow{\sim} \Delta[0] \triangleleft C$, which holds after 4 ~~Lemma 28~~ a) above

4) = RWNEX, which holds by ~~Lemma 28~~ c) above

and 5) = ωES + $\aleph_0$EP which hold by ~~Lemma 28~~ a) and b) above.

This verifies the dual of the strong hypotheses.

The exchange hypothesis of §5, p 136 dualized, cf §5, p 149, follows from c) above

QED

Cahiers Thomason. Bibliothèque MIR

Now Let us check the §5 Th 1.6.1.

Suppose now the $\mathcal{C}$ in addition to having limits there is strongly $\mathcal{M}$-precocious

strongly ω-summable

and satisfies Rich Hypotheses.

§5 Th 1.6.1 Then gives a model structure

$$Cat(\underline{K}, \mathcal{C})$$

Is it Rich ??

Given $F, G : \underline{K} \to \mathcal{C}$.

The internal hom is

$$\underline{Hom}_{Cat(\underline{K}, \mathcal{C})}(F, G) = \int^{\underline{K}^{op} \times \underline{K}} (K, K') \longmapsto \underline{Hom}_{\mathcal{C}}(FK, GK')$$

Note we must take end ~~end~~ here rather than

hoend since I am

$$\left[\underline{Hom}_{Cat(\underline{K}, \mathcal{C})}(F, G)\right]_0 \cong Cat(\underline{K}, \mathcal{C})(F, G)$$

$$\| \int^{end} (K, K') \to \mathcal{C}(FK, GK')$$

$$\| \int^{end} (K, K') \longmapsto \underline{Hom}_{\mathcal{C}}(FK, GK').$$

The end will give an enrichment in Δ^{op}-sets

Cahiers Thomason. Bibliothèque MIR

Best guess for $K\otimes$, $K\triangleleft$ is to form pointwise from $\otimes$, $\triangleleft$ in $\mathcal{E}$.

~~$K\otimes$~~

Let $L : I \to \mathrm{Cat}(\underline{K}, \mathcal{E})$

$L \otimes F : K \longmapsto L \otimes FK$.

$L \triangleleft F : K \longrightarrow L \triangleleft FK$.

This does have the right variances.

$$\begin{aligned}
\mathrm{Nat}(K, \underline{\mathrm{Hom}}(F, G)) &= \mathrm{Nat}(K, \int_{\underline{K}}^{\mathrm{nat}} \underline{\mathrm{Hom}}_{\mathcal{E}}(FL, GL')) \\
&= \int_{\underline{K}}^{\mathrm{nat}} \mathrm{Nat}(K, \underline{\mathrm{Hom}}_{\mathcal{E}}(FL, GL')) \\
&\cong \int_{\underline{K}}^{\mathrm{nat}} \underline{\mathrm{Hom}}_{\mathcal{E}}(FL, K \triangleleft GL') \\
&= \underline{\mathrm{Hom}}_{\mathrm{Cat}}(F, K \triangleleft G)
\end{aligned}$$

and so $\underline{\mathrm{Hom}}_{\mathrm{Cat}}(F \otimes K, G) = \mathrm{Nat}(K, \underline{\mathrm{Hom}}(F, G))$.

But now want to modify definition of fibration

So $G \twoheadrightarrow G'$ is a fibration if

1) $\forall K$ $GK \twoheadrightarrow GK'$ is a fibration (i.e. pointwise)

2) $\forall F \xrightarrow{\sim} F'$ a equicofibration (defined pointwise)

~~$\underline{\mathrm{Hom}}_{\mathrm{Cat}(\underline{K}, \mathcal{E})}(F', G) \longrightarrow \underline{\mathrm{Hom}}_{\mathrm{Cat}(\underline{K}, \mathcal{E})}(F, G) \times \underline{\mathrm{Hom}}_{\mathrm{Cat}(\underline{K}, \mathcal{E})}(F', G')$~~

$\underline{\mathrm{Hom}}(F, G) \times_{\underline{\mathrm{Hom}}(F, G')} \underline{\mathrm{Hom}}(F', G')$

is a K_a fibration (previous def was surjective on $\emptyset \triangleleft [0]$ similarly)

If $F \rightarrowtail F'$, & ~~$L \otimes$~~ $L \rightarrowtail L'$, in

$$L'F \sqcup_{LF} LF' \longrightarrow L'F'$$

is then in pointwise.

Similarly for rest of LE.

Cahiers Thomason. Bibliothèque MIR

So problem is just RE

But given $K \hookrightarrow L$ - f.g. and $G \twoheadrightarrow G'$ in $\underline{Cat}(\underline{E}, \mathcal{C})$

$$L \triangleleft G^{\mathbb{B}} \longrightarrow K \triangleleft G^{\mathbb{B}} \times_{K \triangleleft G} L \triangleleft G'$$

is $F \xrightarrow{\sim} F'$

$$\underline{Hom}(F', L \triangleleft G) \longrightarrow \underline{Hom}(F, L \triangleleft G) \times_{\underline{Hom}(F, K \triangleleft G)} \underline{Hom}(F', K \triangleleft G \times_{K \triangleleft G} L \triangleleft G')$$

$\cong$

$$\underline{Hom}(F, L \triangleleft G) \times \underline{Hom}(F', K \triangleleft G) \times_{\left(\underline{Hom}(F, K \triangleleft G) \times \underline{Hom}(F, L \triangleleft G')\right)^{\underline{Hom}(F, K \triangleleft G)}} \underline{Hom}(F', L \triangleleft G')$$

$$\underline{Hom}(L \otimes F', G) \longrightarrow \underline{Hom}(L \otimes F, G) \times_{\left(\underline{Hom}(K \otimes F, G) \times \underline{Hom}(L \otimes F', G')\right)^{\underline{Hom}(K \otimes F, G)}} \underline{Hom}(K \otimes F', G) \times \underline{Hom}(L \otimes F', G')$$

$\cong$

$$\underline{Hom}(L \otimes F \sqcup_{K \otimes F} K \otimes F', G) \times_{\underline{Hom}(L \otimes F \sqcup_{K \otimes F} K \otimes F', G')} \underline{Hom}(L \otimes F', G')$$

which is a f.l.a.f. $G \longrightarrow G'$,,

$\Leftrightarrow$ s.t. $L \otimes F \sqcup_{K \otimes F} K \otimes F' \xrightarrow{\sim} L \otimes F'$ is ~~cofib~~ equicofinal

$\Leftrightarrow$ $G \longrightarrow G'$ satisfies the condition to be a f.l.a.f.

Cahiers Thomason. Bibliothèque MIR

Moreover, if $K \xrightarrow{\sim} L \rightarrowtail G \twoheadrightarrow G'$ we ⇒ also a

equimorphism, then since $G \twoheadrightarrow G'$ is also a pointwise fibration

is ▽ ⇒ ~~defined~~ coinitial pointwise

$$L \triangleleft G \xrightarrow{\sim} K \triangleleft G \times_{K \triangleleft G} L \triangleleft G'$$ is a pointwise

equimorphism, and hence an equimorphism

So should be able now to lift enriched shan !

$Cat(\mathbb{K}, \mathcal{C})$.

~~Oh~~ Our problem is need to reprove factorization, as / for §5 : 1.6.1-13

definition of f.s.ation has changed.

But should be able to use holim over general

diagram in $\mathcal{C}$ now.

Then since ⇒ ~~$F \rightarrowtail G$~~ $G \twoheadrightarrow G'$ pointwise fib

$$\begin{array}{ccc} G & \xrightarrow{\sim} & G' \times_{\underset{(\mathbb{K}/?)^{\circ}}{\operatorname{holim}} G'} \operatorname{holim}_{(\mathbb{K}/?)^{\circ}} G \longrightarrow \operatorname{holim} G \\ & \searrow & \downarrow \qquad\qquad \downarrow \\ & & G' \longrightarrow \operatorname{holim} G' \end{array}$$

no,
not unless
$\mathcal{C}$ has all
coequalizers !
~~But~~ ~~has~~ ~~grid~~
~~Fonctor~~ ~~not~~ ~~coreflexive~~
~~transfer~~ ~~functor~~
but old trick of
reduction to fibrant
case still works

a: $F \quad F \xrightarrow{\sim} F'$

$$\underline{Hom}\left(F', \operatorname*{holim}_{(\mathbb{K}/?)^{\circ}} G\right) \longrightarrow \underline{Hom}(F', \operatorname{holim} G') \times_{\underline{Hom}(F, \operatorname{holim} G')} \underline{Hom}(F, \operatorname{holim} G)$$

$$= \int^{\text{end}} = \operatorname{holim}_{\mathbb{K}^{\circ} \times \mathbb{K}} \left(\underline{Hom}_{\mathcal{C}}(Fk', Gk') \longrightarrow \underline{Hom}(\cdots \right.$$

so should be OK

Cahiers Thomason. Bibliothèque MIR

We'll also use $\underline{Hom}(F\ \ )$ means pullback

by the $B\Gamma$, th. 4.

Suff. to show

$\forall\ \Delta[n] \in \Delta_f$

$$Mor\left(\Delta[n],\ \underset{\underline{Hom}(F,C)}{\underline{Hom}(F,A) \times \underline{Hom}(F,B)}\right)$$

$$= Mor\left(\Delta[n],\ \underline{Hom}(F,A)\right) \times Mor\left(\Delta[n],\ \underline{Hom}(F,B)\right)$$
$$Mor\left(\Delta[n],\ \underline{Hom}(F,C)\right)$$

ie $Mor_{\mathcal{C}}(F \otimes \Delta[n],\ A) \times Mor_{\mathcal{C}}(F \otimes \Delta[n],\ B)$
$Mor_{\mathcal{C}}(F \otimes \Delta[n],\ C)$

$$= Mor_{\mathcal{C}}\left(F \otimes \Delta[n],\ A \underset{C}{\times} B\right).$$

which is OK by def. of pullback

So back goal,

Lemma: Let $\mathcal{C}$ be a category enriched over Δ^{op}-sets via $\underline{Hom}: \mathcal{C}^{op} \times \mathcal{C} \longrightarrow \Delta^{op}$-sets, and

cf r. 28

suppose for Δ_f = the subcategory of finite simplicial sets

there are functors

$$\Delta_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$$

$$\Delta_f^{op} \times \mathcal{C}^{op} \xrightarrow{\Delta} \mathcal{C}$$

Cahiers Thomason. Bibliothèque MIR

with natural isomorphisms

$$\underline{Hom}(C \otimes K, C') \cong \underline{Map}(K, \underline{Hom}(C, C')) \cong \underline{Hom}(C, K \triangleleft C').$$

Then if $\underline{L} \xrightarrow{F} \mathcal{C}$ is a diagram with a limit $\varprojlim F \in \mathcal{C}$

$$\underline{Hom}(C', \varprojlim F) \longrightarrow \varprojlim \underline{Hom}(C', F)$$ is an isomorphism

in $\underline{\Delta^{op}\text{-sets}}$.

Dually if $\underline{L}^{op} \xrightarrow{F} \mathcal{C}$ is a diagram with a colimit $\varinjlim F \in \mathcal{C}$,

then $$\underline{Hom}(\varinjlim F, C') \longrightarrow \varprojlim \underline{Hom}(F, C')$$

is an isomorphism.

Pf: The "dual statement" is indeed the dual under the duality of §15.

So it suffices to prove the first statement.

As limits are formed pointwise, $Cat(\Delta^{op}, Sets) = \underline{\Delta^{op}Set}$,

it suffices to show $\forall n$ that

$$\left(\underline{Hom}(C', \varprojlim F)\right)_n \longrightarrow \varprojlim \underline{Hom}(C', F)_n$$ is an isomorphism

of sets.

But this map identifies to

$$Map_{\mathcal{C}}(C' \otimes \Delta[n], \varprojlim F) \longrightarrow \varprojlim Map_{\mathcal{C}}(C' \otimes \Delta[n], F)$$

which is an isomorphism by definition of $\varprojlim$.

QED

Cahiers Thomason. Bibliothèque MIR

Prop: Let $\mathcal{C}$ be a left semi-model category (§1, 78)

~~Let λ be an infinite cardinal~~

~~Suppose~~

Suppose arbitrary coequalizers are representable in $\mathcal{C}$.

Suppose $\mathcal{C}$ has coproducts of ~~α~~ objects over index sets of cardinality $< \lambda$. (Note that $\mathcal{C}$ has all colimits over ~~diagram~~ diagrams of cardinality $< \lambda$)

Suppose $\mathcal{C}$ satisfies the other conditions to be strongly λ-coproductive ~~ultra~~ ~~cosummable~~

(§1, p 118, ~~p 120~~) and ultra ω-cosummable (, 44 Solow)

Suppose finally that $\mathcal{C}$ is enriched over Δ^{op}-sets, ~~Suppose there is a functor~~

~~$1_f \times \mathcal{C} \to \mathcal{C}$~~ and has a functor

$1_f \otimes \mathcal{C} \to \mathcal{C}$ of $\mathcal{I}$ a natural iso

$\underline{\mathrm{Hom}}(L \otimes C, C') \cong \underline{\mathrm{Mor}}(L, \underline{\mathrm{Hom}}(C, C'))$

~~which satisfies the~~ and LE (14) hold.

~~axioms for strong left enrichment (this §6 p 24, 26, p 26)~~

Then if $\mathbb{K}$ is a category with $< \lambda$ morphisms:

For any $F : (\Delta/\mathbb{K})^{op} \to \mathcal{C}$

the coend

~~hocofoo~~ $\underset{\Delta/\mathbb{K}}{\mathrm{hocofoo}}\, F = \int^{(\varphi', \varphi) \in \Delta/\mathbb{K} \times (\Delta/\mathbb{K})^{op}} \Delta[\#\varphi'] \otimes F\varphi$

$\cong \int^{(p', p) \in \Delta} \Delta[p'] \otimes \left(\coprod_{\varphi : \mathbb{K} \to \mathbb{K}} F\varphi \right)$

Cahiers Thomason. Bibliothèque MIR

is representable.

If $\eta : F \longrightarrow F'$ is a natural transformation s.t.

$\forall \varphi \in (\Delta/\underline{K})^{op}$ $F\varphi \longrightarrow F'\varphi$ is a cofibration,

then the induced map

$\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, \eta : \underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, F \longrightarrow \underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, F$

is a cofibration.

If each $\eta(\varphi) : F\varphi \longrightarrow F\varphi'$ is an equicofibration,

$\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, \eta$ is a equicofibration.

If $\forall \varphi$ $\eta(\varphi) : F\varphi \longrightarrow F\varphi'$ is an equimorphism

AND if $\forall \varphi$ $F\varphi$, $F\varphi'$ are cofibrant,

then $\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, \eta$ is a equimorphism.

of addendum p 49

haven't assumed functorial factorizat. yet. so don't have this set.

Pf: As $\#(\Delta/\underline{K} \times (\Delta/\underline{K})^{op}) < \aleph_1$, and the coend can be expressed as a colimit over a diagram whose cardinality is then $< \aleph_1$ (in fact the ob. $Sd(\underline{K})$, this is somewhere above.

But also, MacLane, Categories for the Working Mathematician

X 5 Prop 1).

Thus the coend over $\Delta/\underline{K}$ exists, ($\aleph_1$-cocompleteness of $\mathcal{C}$)

The isomorphism with the end over Δ is §3 p 158

As on §§2 §3 p

As on §5 p 6-8, if we filter $\Delta/\underline{K}$ by

sieves $sk_n \Delta/\underline{K} =$ $\varphi : \underline{k} \longrightarrow \underline{K}$ st $\exists$ $q \leq n$ st

$\varphi : \underline{k} \longrightarrow \underline{K}$ factors $\underline{k} \longrightarrow \underline{q} \searrow^{\varphi} \underline{K}$

Cahiers Thomason. Bibliothèque MIR

$(sk_{-1}\,\Delta/K = \emptyset$!

Then

$$\int_{\Delta/K} \Delta[\,] \otimes F \;=\; \varinjlim_{n<\omega} \int_{sk_n \Delta/K} \Delta[\,] \otimes F$$

and $$\int_{sk_{n-1}\Delta/K} \Delta[\,] \otimes F \longrightarrow \int_{sk_n \Delta/K} \Delta[\,] \otimes F$$

f.ts in a pushout square

$$\begin{array}{ccc} \coprod_{\varphi: n \to K \text{ non degenerate}} \dot\Delta[n] \otimes F\varphi & \longrightarrow & \int_{sk_{n-1}\Delta/K} \Delta[\,] \otimes F \\ \downarrow & & \downarrow \\ \coprod_{\varphi: n \to K \text{ non degenerate}} \Delta[n] \otimes F\varphi & \longrightarrow & \int_{sk_n \Delta/K} \Delta[\,] \otimes F \end{array}$$

(but now do not claim left vertical map is cofibration),

This follows from the analysis of factorizat in Δ

(each simplex of Δ/K is degeneracy of a unique non-degenerate simplex)

and the general lemmas §3, 4, 6, and

that $sk_{n-1}(\Delta/K)/\varphi \cong sk_{n-1}\Delta/n$ = colimit of simplices of $\dot\Delta[n]$,

so $\operatorname{colim}_{(sk_{n-1}\Delta/K)/\varphi} (\Delta[\,] \otimes C) \cong (\operatorname{colim}_{sk_{n-1}\Delta/K/\varphi} \Delta) \otimes C \cong \dot\Delta[n]$

Cahiers Thomason. Bibliothèque MIR

using Prop. 3) p. 17 (proof shows part used need so-cat ⊗, not ▷) and ~~Lemma 83 p. 68~~ colim $\Delta[\ ] \cong \Delta[n]$ (ex. §2 p. 5, 68). Also need trivial $\varinjlim \int_{\mathrm{sk}_n \Delta/K} \cong$
similar $\Delta[n]$

Clearly if F takes cofiltrant values, this all reduces to the hocofoo $_{\Delta/K}$ of §5 p. 9.

using Prop. 3) p. 17 (proof shows the part about ⊗ used doesn't need existence of ▷) and colim $\Delta[\ ] \cong \Delta[n]$ (eg §2 p. 5, simplicial $\Delta[n]$
Sus lemma p. 68)

Also need the trivial $\varinjlim_n \int_{\mathrm{sk}_n \Delta/K} = \int_{\cup \mathrm{sk}_n \Delta/K} = \int_{\Delta/K}$

Note if F take cofiltrant values, this hocofoo$_K$ F is indeed the hocofoo$_K$ F of §5 p. 9.

It remains to prove the 2 claims on p. ~~42~~ 41.

~~The last 2 follows from the first~~

Let $\gamma : F \longrightarrow F'$ be a natural transformation
(resp. equicofiltrant)
which for all φ is a cofiltration $\lambda\ \gamma(\varphi) : F\varphi \rightarrowtail F'\varphi$.

~~I claim~~

I need to define:

Cahiers Thomason. Bibliothèque MIR

Def ultra ω-cosummable Let $\mathcal{C}$ be a left semi-model category.

$\mathcal{C}$ is ultra-ω-cosummable if the following condition holds

i) any sequence $C : \omega \longrightarrow \mathcal{C}$ $\quad C_0 \to C_1 \to C_2 \to \cdots$ has a colimit

ii) Let $f_\bullet : C_\bullet \longrightarrow D_\bullet$ be a map of sequences

$$\begin{array}{ccccccc} C_0 & \longrightarrow & C_1 & \longrightarrow & C_2 & \longrightarrow & \cdots \\ {\scriptstyle f_0}\downarrow & & {\scriptstyle f_1}\downarrow & & {\scriptstyle f_2}\downarrow & & \\ D_0 & \longrightarrow & D_1 & \longrightarrow & D_2 & \longrightarrow & \cdots \end{array}$$

Then if

cf 87 v 117

Def: ultra ω-cosummable Let $\mathcal{C}$ be a left semi-model category.

$\mathcal{C}$ is ultra ω-cosummable if the following conditions hold

i) any tower whatsoever in $\mathcal{C}$, $\quad C_0 \to C_1 \to C_2 \to \cdots$ has a colimit $\varinjlim_{\omega} C_i$

ii) Let $f_\bullet : C_\bullet \longrightarrow D_\bullet$ be a map of towers

$$\begin{array}{ccccccc} C_0 & \longrightarrow & C_1 & \longrightarrow & C_2 & \longrightarrow & \cdots \\ {\scriptstyle f_0}\downarrow & & \downarrow{\scriptstyle f_1} & & \downarrow{\scriptstyle f_2} & & \\ D_0 & \longrightarrow & D_1 & \longrightarrow & D_2 & \longrightarrow & \cdots \end{array}$$

suppose that $\forall i : C_i \longrightarrow D_i$ is a cofibration, and that $\forall i : C_{i+1} \underset{C_i}{\sqcup} D_i \longrightarrow D_{i+1}$ is also a cofibration.

(Note the PO exists in $\mathcal{C}$ as $C_i \rightarrowtail D_i$).

Cahiers Thomason. Bibliothèque MIR

Then $\varinjlim_{\omega} C_i \xrightarrow{f} \varinjlim_{\omega} D_i$ is a cofibration.

If in addition each $f_i : C_i \to D_i$ is an equicofibration,

then $\varinjlim_{\omega} C_i \xrightarrow{f} \varinjlim_{\omega} D_i$ is an equicofibration.

—

Now since $\dot\Delta[n] \rightarrowtail \Delta[n]$ is a cofibration in $\mathcal{S}_f$

and $F\varphi \xrightarrow{F'\varphi} F'\varphi$ is a cofibration (resp. equicofibration) in $\mathcal{C}$,

LE gives the

$$\Delta[n]\otimes F\varphi \;\sqcup_{\dot\Delta[n]\otimes F\varphi}\; \dot\Delta[n]\otimes F'\varphi \longrightarrow \Delta[n]\otimes F'\varphi$$

is a cofibration (resp. equicofibration)

By $\mathcal{X}\mathcal{C}$-conduction

$$\coprod_{\varphi:\underline{n}\twoheadrightarrow\underline{k}\ \text{non deg}} \dot\Delta[n]\otimes F\varphi \;\sqcup_{\dot\Delta[n]\otimes F\varphi}\; \dot\Delta[n]\otimes F'\varphi \longrightarrow \coprod_{\varphi\ \text{nd}} \Delta[n]\otimes F'\varphi$$

$\cong$ SII

$$\Big(\coprod_{\varphi\ \text{nd}} \dot\Delta[n]\otimes F\varphi\Big) \sqcup_{\left(\coprod_{\varphi\ \text{n}'} \dot\Delta[n]\otimes F\varphi\right)} \Big(\coprod_{\varphi\ \text{n}'} \dot\Delta[n]\otimes F'\varphi\Big) \longrightarrow \coprod_{\varphi\ \text{nd}} \Delta[n]\otimes F'\varphi$$

is a cofibration (resp. equicofibration).

By ~~LB4~~ we may pushout

I show by induction on n that

$\mathrm{sk}_n \int \Delta[\,]\otimes F \longrightarrow \mathrm{sk}_n \int \Delta[\,]\otimes F'$ is a cofibration (resp. equicofibration)

$\mathrm{sk}_n \Delta/\underline{k}$ $\qquad$ $\mathrm{sk}_n \Delta/\underline{k}$

Cahiers Thomason. Bibliothèque MIR

~~A fact~~

I will prove by induction on n starting at $n=0$

~~that~~

$$\int_{sk_n \Delta/K} \Delta[\,] \otimes F \bigsqcup_{\int_{sk_{n-1}\partial\Delta/K}} \Delta[\,] \otimes F \int_{sk_{n-1}\Delta/K} \Delta[\,] \otimes F' \longrightarrow \int_{sk_n \Delta/K} \Delta[\,] \otimes F$$

~~is a cofibration~~

and $$\int_{sk_n \Delta/K} \Delta[\,] \otimes F \longrightarrow \int_{sk_n \Delta/K} \Delta[\,] \otimes F'$$

are cofibrations (resp. equicofibrations).

~~For start with $n=0$:~~

~~Proof~~

For $n=-1$, $\int_{sk_{-1}} \Delta[\,] \otimes F \longrightarrow \int_{sk_n} \Delta[\,] \otimes F'$ is $\emptyset \xrightarrow{\sim} \emptyset$.

So assume $n \geq 0$, and ~~that~~ by induction hypothesis that I know

$$\int_{sk_{n-1} \Delta/K} \Delta[\,] \otimes F \longrightarrow \int_{sk_{n-1} \Delta/K} \Delta[\,] \otimes F'$$

is a cofibration (resp. ~~trivial~~ equicofibration)

Cahiers Thomason. Bibliothèque MIR

Then LBY gives a pushout

$$\int_{sk_n} \Delta[\,]\otimes F \;\sqcup_{\int_{sk_{n-1}} \Delta[\,]\otimes F} \int_{sk_{n-1}} \Delta[\,]\otimes F' \quad \text{in } \mathcal{C}$$



and the $\int_{sk_n} \Delta[\,]\otimes F \longrightarrow \int_{sk_n} \Delta[\,]\otimes F \sqcup_{\int_{sk_{n-1}} \Delta[\,]\otimes F} \int_{sk_{n-1}} \Delta[\,]\otimes F'$

is a cofibration (resp. equicofibration)

Then using the cofibration (resp. equicofibration) of p45 and the pushout square of p42, I get a pushout square

$$\begin{array}{ccc}
\left(\coprod_{\varphi,\, n\alpha} \dot\Delta[n]\otimes F'_\varphi\right) & \longrightarrow & \coprod_{\varphi,\, n\alpha} \Delta[n]\otimes F_\varphi \\
\downarrow & & \downarrow \\
\left(\coprod_{\varphi} \dot\Delta[n]\otimes F_\varphi\right) & & \\
\downarrow & (PO) & \downarrow \\
\int_{sk_{n-1}} \dot\Delta[\,]\otimes F' & \longrightarrow & \int_{sk_n} \Delta[\,]\otimes F \\
\downarrow & & \\
\int_{sk_{n-1}} \Delta[\,]\otimes F & &
\end{array}$$

By LBY the bottom map is a cofibration (resp. trivial cofibration). This gives the first statement to be proved in the induction step. The second follows

Cahiers Thomason. Bibliothèque MIR

ζ, composition with $\int_{sk_n} \Delta[\,]\otimes F \longrightarrow \int_{sk_n} \Delta[\,]\otimes F \sqcup \int_{sk_{n-1}} \Delta[\,]\otimes F'$

$\int_{sk_{n-1}} \Delta[\,]\otimes F$

~~But~~ This concludes the induction step.

But now, since $\forall n \geq 0$

$$\int_{sk_n \Delta/K} \Delta[\,]\otimes F \rightarrowtail \int_{sk_n \Delta/K} \Delta[\,]\otimes F'$$

ζ

and $\int_{sk_{n+1}\Delta/K} \Delta[\,]\otimes F \sqcup_{\int_{sk_n \Delta/K} \Delta[\,]\otimes F} \int_{sk_n \Delta/K} \Delta[\,]\otimes F' \longrightarrow \int_{sk_{n+1}\Delta/K} \Delta[\,]\otimes F'$

are cofibrations (resp. equicofibrations) ultra ω-cosaturnos.t.

gives that

$$\varinjlim_n \int_{sk_n \Delta/K} \Delta[\,]\otimes F \longrightarrow \varinjlim_n \int_{sk_n \Delta/K} \Delta[\,]\otimes F'$$

$$\int_{\Delta/K} \Delta[\,]\otimes F \qquad \int_{\Delta/K} \Delta[\,]\otimes F'$$

is a cofibration (resp. equicofibration) as required.

QED.

Cahiers Thomason. Bibliothèque MIR

Addendum: If in addition to the hypotheses of Prop 40, $\mathcal{C}$ satisfies left functorial factorization (LFF §3, p 93 cf §1, p 48)

Then, for any ~~weak equivalence~~ natural $\gamma : F \longrightarrow F'$ between functors $\Delta/E \longrightarrow \mathcal{C}$

s.t. 1) $\forall \phi$, $\gamma\phi : F\phi \longrightarrow F'\phi$ is an equimorphism ~~= CAND~~

2) ~~$\forall \phi$, $F\phi$, $F'\phi$ are cofibrant~~, between cofibrant objects

Then $\underset{\Delta/E}{\text{hocolim}}\, \gamma : \underset{\Delta/E}{\text{hocolim}}\, F \longrightarrow \underset{\Delta/E}{\text{hocolim}}\, F'$

is an equimorphism

Pf: Follows from Prop 40 by method of §5, p 12.

Or use §5 Lemma 115 and §5 Prop 32, and Prop 40.

Or note that $F \in \mathrm{Cat}(\Delta/E, \mathcal{C}_{cof})$

the $\underset{\Delta/E}{\text{hocolim}}\, F$ reduces to that of §5 Prop 9, for which the assertion holds. (§5 Prop 9 5), p 11).

Remark: $\mathcal{C} = \Delta^{or}\mathrm{Sets}$ w.r.m. its usual model structure satisfies all the hypotheses of ~~Ex~~ Prop 40.

It also satisfies the dual hypotheses:

to check ~~case~~ ultra ω-summable use method similar to §5 p 176-177

For given tower in Δ^{or}-sets

$$\begin{array}{ccccccc} \to X_n & \longrightarrow & X_{n-1} & \to \cdots & \longrightarrow & X_0 & \to \\ \downarrow f_n & & \downarrow f_{n-1} & & & \downarrow f_0 & \\ \to Y_n & \longrightarrow & Y_{n-1} & \to \cdots & \longrightarrow & Y_0 & \to \end{array}$$

Cahiers Thomason. Bibliothèque MIR

With $f_n : X_n \to Y_n$ and $X_{n+1} \to Y_{n+1} \times_{Y_n} X_n$ Kan fibrations (resp. ~~equivalences~~ weak equivalences and Kan fibrations)

the $\varprojlim X_n \to \varprojlim Y_n$ ~~is~~ is a Kan fibration

(resp. weak equivalence also Kan fibration).

~~If~~ For it suffices to check ~~for~~ the existence of a

fill-in for any diagram ~~of~~ of the form



Where $A \xrightarrow{\sim} B$ is a equivalence (resp. a cofibration).

For the B, universal property of $\varprojlim$ it suffices to find

Compatible fillings :



$\forall n$.

But we construct those ξ_n inductive in n.

For $n = 0$,



such $A \xrightarrow{\sim} B$ (resp. $X_0 \xrightarrow{\sim} Y_0$)

Cahiers Thomason. Bibliothèque MIR

Given $B \longrightarrow X_{n-1}$ we set







But then as $X_n \longrightarrow X_{n-1} \times_{Y_{n-1}} Y_n$ is a fibration,

and either $A \longrightarrow B$ is a weak equivalence,

we have a lift $B \dashrightarrow X_n$ as indicated.

This completes the inductive step of the construction QED

Lemma

Let $\mathcal{C}$ be a category enriched over Δ^{op}-Sets,

with pairs $1_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$ $\quad 1_t^{op} \times \mathcal{C} \xrightarrow{\triangleright} \mathcal{C}$

s.t. $\exists$ natural isomorphisms $\forall L \in 1_f$, $\forall C, C' \in \mathcal{C}$

$$\underline{Hom}(L \otimes C, C') \cong \underline{Mor}(L, \underline{Hom}(C, C')) \cong \underline{Hom}(C, L \triangleright C')$$

Suppose $\mathcal{C}$ is a left model category with functorial factorization (81, p78, 83, p93) and that $\mathcal{C}$ satisfies $)\backslash L$-coproductivity and ω-cosmallness (81, p118, 117)

Suppose that the enrichment satisfies the axiom LE cf p28

Cahiers Thomason. Bibliothèque MIR

a) The $\mathcal{C}$ ~~satisfies~~ satisfies:

the weak hypotheses of §1, 173

the strong hypotheses of §3, 173

the extra hypotheses of §5, 136

~~b)~~ ~~For any $F : \underline{K}$~~

If $\#\underline{K} < \lambda$, for any $F : (\Delta/\underline{K})^{op} \longrightarrow \mathcal{C}_{cof}$

and any $G : (\Delta/\underline{K})^{op} \longrightarrow \mathcal{C}$

there is a natural isomorph

$$\underset{\Delta/\underline{K}}{\mathrm{Hom}}(\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, F,\ G) \cong \underset{\Delta/\underline{K}}{\mathrm{hoend}}\left(K, K' \longmapsto \mathrm{Hom}(FK, GK')\right)$$

cf §5 Thm 140

c) ~~If $\underline{K} \times \underline{K} \xrightarrow{F} \mathcal{C}$ any diagram~~

If $\Phi : \underline{L} \longrightarrow \mathrm{Cat}(\underline{K}, \mathcal{C}_{cof})$ is any

diagram of functors $F_L : \underline{K} \longrightarrow \mathcal{C}_{cof}$

s.t. $\underset{\underline{L}}{\lim} F_L : \underline{K} \longrightarrow \mathcal{C}_{cof}$ exists,

then $\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\left(\underset{\underline{L}}{\lim} F_L\right)$ represents $\underset{\underline{L}}{\lim}\left(L \longrightarrow \underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, F_L\right)$

Pf:

a) follows by the dual to the proof of Lemma 28:

We note this ~~weaker~~ uses only one side of the model structure

Cahiers Thomason. Bibliothèque MIR

5) Given a), 5) follows as in proof of ~~Thm~~ §40 §5 Thm 1.40. Note since the hoend is taken in Δ^{op}-sets, we don't need it to be the ~~the~~ Hom

5) Given a), 5) follows as in the proof of §5 Thm 1.40 5), strictly speaking unneeded hypotheses.

Namely we natural natural isos $\forall n \in \Delta$

$$\underset{\Delta/K}{\underline{\mathrm{Hom}}}(\underset{\Delta/K}{\mathrm{hocolim}}\, F, G)_n = \mathrm{Nat}_{\Delta/K}(\Delta[n] \otimes \underset{\Delta/K}{\mathrm{hocolim}}\, F, G)$$

$\otimes$ is like tensor for Hom ("like" as $\otimes$ defined only on $1_K \in \Delta^{op}/K$)

$$\cong \mathrm{Nat}_{\Delta/K}(\underset{\Delta/K}{\mathrm{hocolim}}\, (\Delta[n] \otimes F), G)$$

§5 Sublemma 7.5

$$\cong \underset{K}{\mathrm{hoend}}\, (K, K' \longmapsto \mathrm{Nat}_{\Delta}(\Delta[n] \otimes FK, GK'))$$

~~FF~~ p 125 5)

$$\cong \underset{K}{\mathrm{hoend}}\, (K, K' \longmapsto \underline{\mathrm{Hom}}(FK, GK')_n)$$

$$\cong (\underset{K}{\mathrm{hoend}}\, (K, K' \longmapsto \underline{\mathrm{Hom}}(FK, GK')))_n$$

limits in Δ^{op}-sets formed pointwise on Δ.

As this iso is natural in $n \in \Delta$, it gives an iso of simplicial sets as required.

(Note: §5 Thm 1.40 asked the ~~Hom(FK, GK')~~ G to take filtrant values so $\underline{\mathrm{Hom}}(FK, GK')$ would be a Kan complex. But we don't need this, hoends (even hocolim's) exist for arbitrary diagrams of Δ^{op}-sets (cf Remark p 49). Then we don't need a CFS, so only need the left model structure of $\mathcal{C}$. Thm means hypotheses of §5 1.40 can be dropped.))

Cahiers Thomason. Bibliothèque MIR

c). For any $X \in \mathcal{C}$

$$\mathrm{Mor}_{\mathcal{C}}\left(\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\left(\underset{\underline{K}}{\mathrm{lim}}\, F_L\right),\ X\right)$$

$$= \underset{\underline{K}}{\mathrm{hoend}}\left(K, K' \right)$$

cf §5 r140,

Prop. Let $\mathcal{C}$ be a category enriched over Δ^{op}-sets with $\mathcal{A}_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$ and $\mathcal{A}_f^{op} \times \mathcal{C} \xrightarrow{\triangleright} \mathcal{C}$ s.t. $\exists$ natural isos $\quad \forall\ L \in \mathcal{A}_f,\ C, C' \in \mathcal{C}$

$$\underline{\mathrm{Hom}}(L \otimes C, C') \cong \mathrm{Mor}(L, \underline{\mathrm{Hom}}(C, C')) \cong \underline{\mathrm{Hom}}(C, L \triangleright C')$$

Suppose
$\mathcal{C}$ is a left model category with functorial factorization (§1 r48, §3 r93) which satisfies $\mathcal{M}$-coproducts and $\mathcal{M}$-cosmashability. (§1 r118, 117).

Suppose further the enrichment satisfies LE of r28.

Suppose $\#\,\mathrm{ob}\,\underline{K} < \mathcal{M}$.

Then

a) For any $F : (\Delta/\underline{K})^{op} \longrightarrow \mathcal{C}_{cof}$

and any $C' \in \mathcal{C}$

notation
cf §5 r154-157

$$\underline{\mathrm{Hom}}\left(\underset{\Delta/\underline{K}}{\mathrm{hocofoo}}\, F,\ C'\right) \cong \underset{(\Delta/\underline{K})^{op}}{\mathrm{hofoo}}\left(\varphi \longmapsto \underline{\mathrm{Hom}}(F\varphi, C')\right)$$

b) For any $F : \underline{K} \longrightarrow \mathcal{C}_{cof}$

and any $G : \underline{K} \longrightarrow \mathcal{C}$

there is a natural isomorphism

Cahiers Thomason. Bibliothèque MIR

$$\int^{K \in \mathcal{K}^{op}} (K, K') \longmapsto \underline{\mathrm{Hom}}_{\mathcal{C}}\Big(\operatorname*{hocolim}_{K \in \mathcal{K}/K} F,\ GK'\Big)$$

$$\cong \operatorname*{hoend}_{K \in \mathcal{K}} \big((K, K') \longmapsto \underline{\mathrm{Hom}}_{\mathcal{C}}(FK, GK')\big)$$

[hoend a special case of hoSaun

c) If $\Phi : \mathcal{L} \longrightarrow \mathrm{Cat}\big((\Delta/\mathcal{K})^{op}, \mathcal{C}_{cof}\big)$

is any diagram of functors s.t.

$\operatorname*{hocolim}_{L \in \mathcal{L}} \Phi L$ exists as a functor $(\Delta/\mathcal{K})^{op} \longrightarrow \mathcal{C}_{cof}$

(not it is assumed to take cofibrant values!).

Then $\operatorname*{hocofoo}_{\Delta/\mathcal{K}} \big(\operatorname*{hocolim}_{L \in \mathcal{L}} \Phi L\big)$ represents

$$\operatorname*{hocolim}_{L \in \mathcal{L}} \big(L \longmapsto \operatorname*{hocofoo}_{\Delta/\mathcal{K}} \Phi L\big).$$

Pf:

Notes a) By Lemma 51, $\mathcal{C}$ satisfies the weak hypotheses, strong hypotheses, and extra hypotheses. Thus $\operatorname*{hocofoo}_{\Delta/\mathcal{K}} F$ exists by 88 Prop 9.

$\operatorname*{hofoo}_{(\Delta/\mathcal{K})^{op}}$ exists as Δ^{op}-set for any functor, without need to assume fibrant values. (Prop 140 + Remark 141)

b) So both sides exist.

Then ~~there~~ $\exists$ natural isomorphism in $F, \mathcal{K}, \mathcal{C}'$ and $n \in \Delta$

$$\underline{\mathrm{Hom}}\Big(\operatorname*{hocofoo}_{\Delta/\mathcal{K}} F, \mathcal{C}'\Big)_n \cong \mathrm{Mor}_{\mathcal{C}}\big(\Delta[n] \otimes \operatorname*{hocofoo}_{\Delta/\mathcal{K}} F, \mathcal{C}'\big)$$

$\otimes$ the partial cotensor

$$\cong \mathrm{Mor}_{\mathcal{C}}\Big(\operatorname*{hocof}_{\Delta/\mathcal{K}} \Big)$$

Cahiers Thomason. Bibliothèque MIR

$$\simeq \mathrm{Map}_{\mathcal{C}}\Big(\operatorname*{hocofoo}_{\Delta/\mathcal{K}} \Delta[n]\otimes F,\ C'\Big)$$

[§5 Lemma 7.5

$$\simeq \mathrm{Map}_{\mathcal{C}}\Big(\int^{\Delta/\mathcal{K}\times(\Delta/\mathcal{K})^{op}} \Delta[\,]\otimes(\Delta[n]\otimes F),\ C'\Big)$$

[def of hocofoo

$$\simeq \int_{(\Delta/\mathcal{K})^{op}\times\Delta/\mathcal{K}}^{end} \mathrm{Map}_{\mathcal{C}}\big(\Delta[\,]\otimes(\Delta[n]\otimes F),\ C'\big)$$

[& universal property of coend

$$\simeq \int_{(\Delta/\mathcal{K})^{op}\times\Delta/\mathcal{K}}^{end} \mathrm{Map}_{\mathcal{C}}\big(\Delta[\,]\otimes(\Delta[\,]\times\Delta[n])\otimes F,\ C'\big)$$

{ enriched + part isomorphism

$$\simeq \int_{(\Delta/\mathcal{K})^{op}\times\Delta/\mathcal{K}}^{end} \mathrm{Map}_{\mathcal{C}}\big(\Delta[n]\otimes(\Delta[\,]\otimes F),\ C'\big)$$

$$\simeq \int_{(\Delta/\mathcal{K})^{op}\times\Delta/\mathcal{K}}^{end} \underline{\mathrm{Hom}}\big(\Delta[\,]\otimes F,\ C'\big)_n$$

$$\simeq \left[\int_{(\Delta/\mathcal{K})^{op}\times\Delta/\mathcal{K}}^{end} \underline{\mathrm{Hom}}\big(\Delta[\,]\otimes F,\ C'\big)\right]_n$$

ends pturise in Δ^{op}-sets

$$\simeq \left[\int_{(\Delta/\mathcal{K})^{op}\times\Delta/\mathcal{K}}^{end} \underline{\mathrm{Map}}\big(\Delta[\,],\ \underline{\mathrm{Hom}}(F, C')\big)\right]_n$$

$$= \Big[\operatorname*{hofoo}_{(\Delta/\mathcal{K})^{op}} \underline{\mathrm{Hom}}(F, C')\Big]_n$$

{ def of hofoo of §5, 154, 157 and §Thm 8.6, 170.

As natural in n, this gives a natural iso of simplicial sets

$$\underline{\mathrm{Hom}}\Big(\operatorname*{hocofoo}_{\Delta/\mathcal{K}} F,\ C'\Big) \simeq \operatorname*{hofoo}_{(\Delta/\mathcal{K})^{op}} \underline{\mathrm{Hom}}(F, C')$$

Cahiers Thomason. Bibliothèque MIR

This proves a)

b) Ends exist in the complete category Δ^n-sets, as do hoends of arbitrary functors (cf Prop 40 + Remark 79).

~~end~~

There is a isomorphism natural in $\underline{\underline{K}}, K, K', F, G$ and n.

$$\left[\int_{\underline{\underline{K}}}^{\text{end}} (K,K') \longmapsto \underline{\text{Hom}}_{\mathcal{E}} \left(\operatorname*{hocolim}_{\underline{\underline{K}}/K} F, GK' \right) \right]_n \cong \int_{\underline{\underline{K}}}^{\text{end}} (K,K') \longmapsto \left[\underline{\text{Hom}}_{\mathcal{E}} \left(\operatorname*{hocolim}_{\underline{\underline{K}}/K} F, GK' \right) \right]_n$$

$$\cong \int_{\underline{\underline{K}}}^{\text{end}} (K,K') \longmapsto \underline{\text{Mor}}_{\mathcal{E}} \left(\Delta[n] \otimes \operatorname*{hocolim}_{\underline{\underline{K}}/K} F, GK' \right)$$

$$\cong \int_{\underline{\underline{K}}}^{\text{end}} (K,K') \longmapsto \text{Mor}_{\mathcal{E}} \left(\operatorname*{hocolim}_{\underline{\underline{K}}/K} F \otimes \Delta[n], GK' \right)$$

[suslemm 85 r 75 / cond simpl.]

$$\cong \left[\operatorname*{hoend}_{\underline{\underline{K}}} (K,K' \longmapsto \underline{\text{Hom}}_{\mathcal{E}}(FK, GK') \right]_n$$

$$\cong \text{Mor}_{\text{Cat}(\Delta/\underline{\underline{K}}, \mathcal{E})} \left(\operatorname*{hocolim}_{\underline{\underline{K}}/?} (F \otimes \Delta[?]), G \right)$$

[definition of natural transformation]

$$\cong \text{Mor}_{\text{Cat}((\Delta/\underline{\underline{K}})^{\text{op}}, \mathcal{E})} \left(\operatorname*{hocolim}_{\underline{\underline{K}}/?} (F \otimes \Delta[?]), G \right)$$

$$\cong \text{Mor}_{\text{Cat}(\underline{\underline{K}}, \mathcal{E})} \left(\operatorname*{hocolim}_{\underline{\underline{K}}/?} F \otimes \Delta[?], G \right)$$

[85 r 127]

$$\cong \left[\operatorname*{hoend}_{\underline{\underline{K}}^{\text{op}}} (K,K' \longmapsto \underline{\text{Hom}}_{\mathcal{E}}(F \otimes \Delta[n], G) \right]_0$$

$$\cong \left[\operatorname*{hoend}_{\underline{\underline{K}}^{\text{op}}} (K,K') \longmapsto \text{Mor}(\Delta[n], \underline{\text{Hom}}_{\mathcal{E}}(F,G)) \right]_0$$

Cahiers Thomason. Bibliothèque MIR

$$\cong \underset{\Delta^{op}\to Set}{Map}\left(\Delta[n],\ \left(\underset{\mathbb{K}}{hoend}\ K,K' \longrightarrow \underline{Hom}_e(FK, GK')\right)\right)$$

But in $\Delta^{op}Set$ $\underline{Map}(\Delta[n],\ )$ preserves all limits & colimits, hence all ends. Thus

$$\underset{\mathbb{K}}{hoend}\ (K,K' \longrightarrow \underline{Map}(\Delta[n],\ \underline{Hom}(FK, GK'))$$

cf §7 Statement 5

$$= \int_{(\Delta/\mathbb{K})^{op}} \underline{Map}(\Delta[\ ],$$

$$\int_{(\Delta/\mathbb{K})^{op}} (\varphi',\varphi) \longmapsto \underline{Map}(\Delta[\varphi'],\ \underline{Map}(\Delta[n],\ \underline{Hom}(F\varphi 0,\ G\varphi(mor)))$$

$$\cong \int^{op}_{(\Delta/\mathbb{K})^{op}} (\varphi',\varphi) \longmapsto \underline{Map}(\Delta[\varphi'] \times \Delta[n],\ \underline{Hom}(F\varphi 0,\ G\varphi(mor)))$$

$$\cong \int^{end}_{(\Delta/\mathbb{K})^{op}} (\varphi',\varphi) \longmapsto \underline{Map}(\Delta[n],\ \underline{Map}(\Delta[\varphi'],\ \underline{Hom}(F\varphi 0,\ G\varphi(mor))))$$

$$\cong \underline{Map}(\Delta[n],\ \int^{end}_{(\Delta/\mathbb{K})^{op}} (\varphi',\varphi) \longmapsto \underline{Map}(\Delta[\varphi'],\ \underline{Hom}(F\varphi 0,\ G\varphi(mor))))$$

$$= \underline{Map}(\Delta[n],\ \underset{\mathbb{K}}{hoend}\ \underline{Hom}_e(F,G))$$

So $$\left[\underset{\mathbb{K}^{op}}{hoend}\ (K,K') \longmapsto \underline{Map}(\Delta[n],\ \underline{Hom}(FK, GK'))\right]_0$$

$$\cong \left[\underline{Map}(\Delta[n],\ \underset{\mathbb{K}^{op}}{hoend}\ \underline{Hom}_e(F,G))\right]_0$$

$$= \left[\underset{\mathbb{K}}{hoend}\ \underline{Hom}_{\mathcal{C}^{op}}(F,G)\right]_n.$$

Cahiers Thomason. Bibliothèque MIR

As this isomorphism is natural in a, it gives us the desired natural isomorphism of simplicial sets.

$$\int^{a \in \mathbb{\Delta}} (K, K') \longmapsto \underline{\mathrm{Hom}}_e(\operatorname*{hocolim}_{\mathbb{\Delta}/K} F,\ G)$$

$$\cong \operatorname*{holim}_{\mathbb{\Delta}} (K, K' \longmapsto \underline{\mathrm{Hom}}_e(FK, GK')).$$

This proves 5).

4) For any $C \in \mathcal{C}$

$$\mathrm{Mor}_{\mathcal{C}}\Big(\operatorname*{hocofoo}_{\mathbb{\Delta}/K} F, C\Big)$$

$$\mathrm{Mor}_{\mathcal{C}}\Big(\operatorname*{hocofoo}_{\mathbb{\Delta}/\mathbb{K}} \Phi L,\ C\Big)$$

$$= \Big[\operatorname*{hofoo}_{(\mathbb{\Delta}/\mathbb{K})^{op}} \big(\varphi \longmapsto \underline{\mathrm{Hom}}_e\big(\operatorname*{lim}_{\mathbb{\Delta}} \Phi L(\varphi),\ C\big)\big)\Big]_0$$

$$= \operatorname*{hofoo}_{\mathbb{\Delta}/\mathbb{K}^{op}} \big(\varphi \longmapsto \operatorname*{lim}_{\mathbb{\Delta}^{op}} \mathrm{Hom}$$

c) $\operatorname*{hofoo}_{\mathbb{\Delta}}$ $\operatorname*{hocofoo}_{\mathbb{\Delta}/\mathbb{K}}\big(\operatorname*{lim}_{\mathbb{\Delta}} \Phi L\big)$, $\operatorname*{hocofoo}_{\mathbb{\Delta}/\mathbb{K}} \Phi L$ exist.

$\operatorname*{lim}_{\mathbb{\Delta}} \Phi L$, $\Phi L : \mathbb{\Delta}/\mathbb{K} \longrightarrow$ $\mathcal{C}$ cof.

Cahiers Thomason. Bibliothèque MIR

Then

$$\underset{\Delta/\mathbb{E}}{\mathrm{hocolim}}\ \left(\varinjlim_{\mathbb{E}} \Phi L\right)$$

$$= \int^{\Delta/\mathbb{E}} (\varphi', \varphi) \longmapsto \Delta[\#\varphi'] \otimes \left(\varinjlim_{\mathbb{E}} \Phi L\right)$$

$$\simeq \int^{\Delta/\mathbb{E}} (\varphi', \varphi) \longmapsto \varinjlim_{\mathbb{E}} \left(\Delta[\#\varphi'] \otimes \Phi L\right)$$

$$\simeq \varinjlim_{\mathbb{E}} \int^{\Delta/\mathbb{E}} (\varphi', \varphi) \longmapsto \Delta[\#\varphi'] \otimes \Phi L$$

$$= \varinjlim_{\mathbb{E}} \left(L \longmapsto \underset{\Delta/\mathbb{E}}{\mathrm{hocolim}}\ \Phi L\right).$$

QED.

argument
$p(6\ 4)$ maybe ~~it~~ one-sided
~~we do~~
ours needs a left model structure, but enrichment on both sides.

colimits commute

~~[includes representables, etc.]~~ !]
~~of such~~

[includes representables, etc. by statement]

Cahiers Thomason. Bibliothèque MIR

Thm: Let $(\mathcal{C}, \mathrm{cof}\mathcal{C}, \mathrm{equi}\mathcal{C}, \mathrm{fis}\mathcal{C})$ satisfy the basic model axioms (78 156–157) B1–B6, and functorial factorization F1, F'1 (78, 158) (These imply B7. Given richness, will follow from C1 of 79, 172).

Suppose $\mathcal{C}$ is enriched over Δ^{or}-sets, and that there are "partial" tensor and cotensor

$$1_f \times \mathcal{C} \xrightarrow{\otimes} \mathcal{C}$$
$$1_f^{\mathrm{op}} \times \mathcal{C} \xrightarrow{\triangleright} \mathcal{C}$$

($1_f \in \Delta^{\mathrm{or}}\text{-sets}$ = full subcategory of finite Δ^{or}-sets.

s.t.

$$\mathrm{Hom}(X \otimes C, C') \cong \mathrm{Map}(X, \mathrm{Hom}_{\mathcal{C}}(C, C')) \cong \mathrm{Hom}_{\mathcal{C}}(C, X \triangleright C')$$

Suppose axioms RE and LE hold (p. 14, 15).
Suppose that $\mathcal{C}$ satisfies $\aleph_1$-productivity and ω-summasivity (81, 118, 117).

Let $\mathbb{K}$ be a category with $\#\mathrm{mor}\,\mathbb{K} < \aleph_1$.

Used $\mathbb{K}^{\mathrm{or}}$ in 85 p. 161 but no where else.

Then $\mathrm{Cat}(\mathbb{K}, \mathcal{C})$ has a basic model structure with functorial factorization, $\mathcal{C}$ is enriched over Δ^{or}-sets with $\otimes$, $\triangleright$ s.t. LE + RE hold, and $\mathrm{Cat}(\mathbb{K}, \mathcal{C})$ satisfies $\aleph_1$-productivity and ω-summasivity.

This structure is given by

$\gamma: F \to G$ a natural transformation is

i) a cofibration if $\forall K \in \mathbb{K}$, $\gamma K: FK \to GK$ is a cofibration in $\mathcal{C}$

ii) an equimorphism if $\forall K \in \mathbb{K}$, $\gamma K: FK \to GK$ is an equimorphism in $\mathcal{C}$

Cahiers Thomason. Bibliothèque MIR

iii) a fibration if $\forall K \in \underline{K}$ $FK \longrightarrow GK$ is a fibration in $\mathcal{C}$ and for all equicofibration $A \xrightarrow{\sim} B$ in $Cat(\underline{K}, \mathcal{C})$

$$\underline{Hom}_{Cat(\underline{K},\mathcal{C})}(B, F) \longrightarrow \underline{Hom}_{Cat(\underline{K},\mathcal{C})}(A, F) \times \underline{Hom}(B, G)$$
$$\underline{Hom}(A, G)$$

is a Kan fibration and weak equivalence of simplicial sets.

a) For $X \in \mathcal{S}_f$ and $F : \underline{K} \longrightarrow \mathcal{C}$

$X \otimes F : \underline{K} \longrightarrow \mathcal{C}$ sends $K \longmapsto X \otimes FK$.

$X \triangleleft F$ ~~sends~~ $: \underline{K} \longrightarrow \mathcal{C}$ sends $K \longmapsto X \triangleleft FK$.

b) For $F, G \in Cat(\underline{K}, \mathcal{C})$.

$$\underline{Hom}_{Cat(\underline{K},\mathcal{C})}(F, G) = \int_{\underline{K}}^{end} (K, K') \longmapsto \underline{Hom}_{\mathcal{C}}(FK, GK')$$

c) A natural transformation $\gamma : F \longrightarrow G$ of functors $\underline{K} \longrightarrow \mathcal{C}$ is a cofibration if it is pointwise a cofibration in $\mathcal{C}$ ($\forall K \in \underline{K}$, $\gamma K : FK \longrightarrow GK$ in $\mathcal{C}$).

~~If~~ γ is a equimorphism if it is so pointwise ($\forall K \in \underline{K}$, $\gamma K : FK \xrightarrow{\sim} GK$ is in $\mathcal{C}$.

γ is a fibration if it is pointwise a fibration and for all diagrams

Cahiers Thomason. Bibliothèque MIR



where $A \rightarrowtail B$ is a trivial cofibration, then

$\exists$ a l.s.t.

Pf: By the dual to p 51, $\mathcal{C}$ satisfies the dual weak hypotheses (§1 p 143 dualized), the dual strong hypotheses (§3 p 173, dualized) and the dual extra hypotheses of §3 (§5, 136). These give JXCEP (§5, 149) & RWN (§5, 149), since these are among the dual strong hypotheses.

By §5 Thm p 161, $Cat(\underline{K}^{2n}, \mathcal{C})$ with the described structure is a basic model category with functorial factorization, which is JXC-productive and ω-summable.

Moreover $\underline{Hom}(F,G) = \int^{(K,K') \in \underline{K}^{2n}} (K,K') \longmapsto \underline{Hom}_{\mathcal{C}}(FK, GK')$

is easily seen to give an enrichment over Δ^{cc}-set.

(Note $\left[\underline{Hom}(F,G)\right]_0 = \left[\int^{(K,K') \in \underline{K}^{2n}} (K,K') \longmapsto \underline{Hom}_{\mathcal{C}}(FK, GK')\right]_0$

$= \int^{(K,K') \in \underline{K}^{2n+1}} \left[(K,K') \longmapsto \underline{Hom}_{\mathcal{C}}(FK, GK')\right]$

$= \int^{(K,K') \in \underline{K}^{2n}} (K,K') \longmapsto Mor_{\mathcal{C}}(FK, GK'))$

$= Mor_{Cat(\underline{K},\mathcal{C})}(F,G).$

Cahiers Thomason. Bibliothèque MIR

Moreover [$X \subset I_k$

$$\underline{\mathrm{Hom}}(X \otimes F, G)$$

$$= \int^{\mathrm{end}} (K, K') \vdash \underline{\mathrm{Hom}}_e(X \otimes FK, GK')$$

$$= \int^{\mathrm{end}} (K, K') \vdash \underline{\mathrm{Mor}}(X, \underline{\mathrm{Hom}}_e(FK, GK'))$$

$$\cong \underline{\mathrm{Mor}}(X, \int^{\mathrm{end}} (K, K') \vdash \underline{\mathrm{Hom}}_e(FK, GK'))$$

$$\cong \underline{\mathrm{Mor}}(X, \underline{\mathrm{Hom}}(F, G))$$

$$\cong \int^{\mathrm{end}} (K, K') \vdash \underline{\mathrm{Mor}}(X, \underline{\mathrm{Hom}}(FK, GK'))$$

$$= \int^{\mathrm{end}} (K, K') \vdash \underline{\mathrm{Hom}}_e(FK, X \triangleright GK'))$$

$$\cong \underline{\mathrm{Hom}}(F, X \triangleright G).$$

It remains only to verify the axiom LE or RE

Actually both.

But given $X \longrightarrow Y$ in I_f a $F \longrightarrow G$. Cat($\underline{\mathrm{Ens}}_e$)

$Y \otimes F \sqcup_{X \otimes F} X \otimes G \longrightarrow Y \otimes G$ is indeed representable

by a colimitation, since this may be determined

pointwise, and $\forall K \in I_k$

$Y \otimes FK \sqcup_{X \otimes FK} X \otimes GK \longrightarrow Y \otimes GK$ is represented

Cahiers Thomason. Bibliothèque MIR

by a cofibration in $\mathcal{C}$ by $LE \subset \mathcal{C}$.

Now to check RE.

Let X

If either $X \rightarrowtail Y$ or $F \rightarrowtail G$ is an equivalence,

so is $Y \otimes F \amalg_{X \otimes F} X \otimes G \longrightarrow Y \otimes G$, by LE in $\mathcal{C}$,

this condition being verifiable pointwise.

Now $\subset$ RE.

Let $X \rightarrowtail Y$ be in and $F \longrightarrow G$ be a fibration.

Then $X \triangleright F \longrightarrow X \triangleright G$ ×

The $Y \triangleright F \longrightarrow X \triangleright F \times_{X \triangleright X} Y \triangleright G$

is representable by a pointwise fibration, which is

moreover an equivalence if either $X \xrightarrow{\sim} Y$ or $F \xrightarrow{\sim} G$.

It remains only to show $\subset$ $X \rightarrowtail Y$ or $F \twoheadrightarrow G$

that the pointwise fibration

$$Y \triangleright F \longrightarrow X \triangleright F \times_{X \triangleright G} Y \triangleright G \quad \text{is a fibration,}$$

ie that it also satisfies the lifting property of

the definition of fibration.

But let $A \xrightarrow{\sim} B$ be an equicofibration,

Cahiers Thomason. Bibliothèque MIR

Note there is an adjunct $\Gamma, X \in I_\Sigma$

$$X \otimes \dashv X \triangleright$$

For $\underline{Hom}(X \otimes A, F) \cong \underline{Mor}(X, \underline{Hom}(A, F)) \cong \underline{Hom}(A, X \triangleright F)$

so taking ΣI_0

$$Mor(X \otimes A, F) \cong Mor(A, X \triangleright F).$$

Then



correspond bijections to



Since $X \in (I_\Sigma) = (I_\Sigma)$ cof, $X \otimes A \to X \otimes B$, our these

diagrams correspond bijections to



By LF on $X \otimes B \underset{X \otimes A}{\sqcup} Y \otimes A \xrightarrow{\sim} Y \otimes B$ is a equiv. by 2 out of 3 since $X \otimes A \xrightarrow{\sim} B$ is.

Cahiers Thomason. Bibliothèque MIR

But then by definition of fibration a dotted arrow exists in the bottom square.

Under the correspondence this gives the dotted arrow exists in the upper square, as required to show the lifting property. QED.

Let's see how this if the new model category fixes the complaint of Quillen in Homological Algebra II 6.3, II 4.1, 4.2

Let $\underline{A}$ be a category with finite limits and finite colimits (what Quillen means: closed under finite limits)

Suppose $\forall X \in \underline{A}$ $\exists$ a projective object $P \in \underline{A}$ and an effective epimorphism $P \longrightarrow X$ (Quillen: $\underline{A}$ has enough projectives)

Let $s\underline{A}$ be the category of simplicial objects over $\underline{A}$

It is enriched over Δ^{op}-sets with colimits and tensors over sets by Quillen Homological Algebra, II 1.7-1.8

Say a morphism $A. \xrightarrow{f} B.$ is

~~a fibration~~ if

1) a fibration if $\forall P \in \underline{A}$ projective

$\underline{Hom}(P, A.) \longrightarrow \underline{Hom}(P, B.)$ is a Kan fibration

2) a equivalence if $\forall P \in \underline{A}$ projective

$\underline{Hom}(P, A) \longrightarrow \underline{Hom}(P, B)$ is a weak equivalence of Δ^{op}-sets.

Cahiers Thomason. Bibliothèque MIR

3) a cofibration of $A. \longrightarrow B.$ is a monomorph... with $B.V_n \quad B_n/A_n$ a projective object.

Check this gives a model structure satisfying the enriched hypotheses.

Check Basic model axioms 78 v156-157

B1 clear

B2 clear.

B3 clear

B4. Given

$$\begin{array}{ccc} A. & \longrightarrow & B. \\ \downarrow & & \\ C. & & \end{array}$$

The pushout $C. \underset{A.}{\cup} B.$ exists as $\underline{A}$, hence $s\underline{A}$ has all finite colimits.

The cofibration $C. \underset{A.}{\cup} B./C. \cong B./A.$ and so is projective in each degree.

To conclude $C. \longrightarrow C. \underset{A.}{\cup} B.$ is a cofibration, it remains to check it is mono.

Argh! No reason why it should be mono.

Cahiers Thomason. Bibliothèque MIR

OK Choose definition of cofibration.

Go back & Quillen's

$A \longrightarrow B$ is a cofibration iff it has the left lifting property for the equifibrations.

Restart.

B1, B2, B3 still ok

B4: ~~Clear from definition~~

half LB3: sA has all finite colimits as A does.

So PO's exist.

Clear from definition that PO of a cofibration is a cofibration. Lifting property is preserved.

Sum

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & D & \downarrow \\ C & \longrightarrow & D \end{array}$$

as $A \xrightarrow{\sim} B$.

Why is $C \xrightarrow{\sim} D$?

- $\mathrm{Hom}(P, C) \xrightarrow{\sim} \mathrm{Hom}(P, D)$.

No good reason. LB3 shd

RB4 ok. Pullbacks exist, as $\mathrm{Hom}(P, \ )$ converts pullbacks to pullbacks. As $\Delta^\circ \mathrm{Set}$ is a proper model cat, the sim. RB4.

Cahiers Thomason. Bibliothèque MIR

B5 trivial

B6 has fill in & def.

B7 Quillen II 4.5 Pr 3



This gives half of B7. Put off other half.

B6 has admiss fill in & def.

Gm

would like to use trick of §5 p165, dualn







have on RB4

half of B7 we [illegible] su





Cahiers Thomason. Bibliothèque MIR

we need the other functors ($\tilde{\longrightarrow}$, $\twoheadrightarrow$),

No so back to first idea

$A. \longrightarrow B.$ a cofibration iff

$\forall n$ $A_n \longrightarrow B_n$ " mono with B_n/A_n projective

Better $A. \longrightarrow B.$ a cof. iff

$\forall n$ $B_n \cong A_n \underset{\emptyset}{\cup} P_n$ with P_n projective

(equivalent in case A additive, but not in general

Now easy that the [illegible] of base change preserves cofibs

$$\begin{array}{ccc} A. & \longrightarrow & B. \\ \downarrow & & \downarrow \\ C. & \longrightarrow & C \underset{A}{\cup} B \end{array}$$

$C. \longrightarrow C \underset{A}{\cup} B.$ " $C_n \longrightarrow C_n \underset{\emptyset}{\cup} P_n$!!

If ~~$A. \tilde{\longrightarrow} B.$~~ "

But properties still a ~~=~~ a problem

Nice $A_n \longrightarrow A_n \cup P_n$ " $B.$ . . . $A. \tilde{\longrightarrow} B.$

th $P_n = B./A.$ " $\simeq 0.$

$0 \leftarrow \mathrm{Hom}(Q, A_n) \longrightarrow \mathrm{Hom}(Q, A_n \underset{\emptyset}{\cup} P_n) \leftarrow \mathrm{Hom}(Q, P_n) \leftarrow 0$

NOT "EXACT" in non-additive case

Cahiers Thomason. Bibliothèque MIR

Somewhat VAGUE
IDEA

Better solution to Quillen's complaint argum

$s\underline{A}$: Let

Consider $\underline{A}$ as a site with the canonical topology

Note P preserves gives on with trivial torsors

give equivalent topos.

Now $s\underline{A} \in \Delta^{op}\Delta^{\wedge} \hookrightarrow \Delta^{op}P^{\wedge} \simeq \Delta^{op}$ (sheaves on A)

which is a topos by Joyal.

Quillen structure is restriction of structure of Joyal.

Only Problem is to check that shom $s\underline{A}$ is

closed under the factorization which exists Δ^{op} (sheaves on $\underline{A}$)

—

Other problem

$\underline{\underline{C}}$ a site $\mathcal{C}$ model category satisfying rich hypothesis

Cat$\underline{\underline{C}}$ $\#\underline{\underline{C}} < \lambda$,

Sum $\mathcal{C}$ as λ-products + ω-sums

$\mathcal{C}$ as λ-colimits + λ-limits !

Then $Cat(\underline{\underline{C}}^{op}, \mathcal{C})$ has model structure

satisfying rich hypotheses.

$Sheaves(\underline{\underline{C}}, \mathcal{C}) \subseteq Cat(\underline{\underline{C}}^{op}, \mathcal{C})$

F is $\{U_i \to U\}$ a cover

$FU \to \prod FU_i \rightrightarrows \prod F(U_i \times_U U_j)$

Cahiers Thomason. Bibliothèque MIR

Can we make a sheafification functor, and lift model structure to Sheaves.

Maybe we

Suppose $\mathcal{A} \underset{i}{\overset{sh}{\rightleftarrows}} \mathcal{C}$

with $sh \dashv i$ $\quad sh \circ i \xrightarrow{\cong} Id$ so i fully faithful

$\mathcal{C}$ has a ~~enriched~~ model structure satisfying the ~~right~~ axioms

however.

Can we lift $\mathcal{A}$ to $\mathcal{C}$?

~~$\underline{Hom}_{\mathcal{A}}(A, A') = \underline{Hom}_{\mathcal{C}}(iA, iA')$~~ ?

~~$K \otimes_{\mathcal{A}} A = sh(K \otimes_{\mathcal{C}} iA)$~~

$K \otimes_{\mathcal{A}} A = sh(K \otimes_{\mathcal{C}} iA)$

$K \pitchfork_{\mathcal{A}} A = sh(K \pitchfork_{\mathcal{C}} A)$

Then $\underline{Hom}_{\mathcal{A}}(sh(K \otimes_{\mathcal{C}} iA), A') = \underline{Hom}_{\mathcal{C}}(i \circ sh(K \otimes_{\mathcal{C}} iA), iA')$

$\underline{Hom}_{\mathcal{A}}(shC, A') = \underline{Hom}_{\mathcal{C}}(ish\, C, iA')$

$\simeq \underline{Hom}_{\mathcal{C}}(C, iA')$?

No. so $K \otimes_{\mathcal{A}} A = sh(K \otimes_{\mathcal{C}} iA)$.

The $\underline{Hom}$ def $\underline{Hom}_{\mathcal{A}}$ is $\underline{Hom}_{\mathcal{A}}(A, A')_n = Mor_{\mathcal{A}}(\Delta[n] \otimes A, A')$

Cahiers Thomason. Bibliothèque MIR

$$= \underline{Mor}_e(i'(\Delta[n] \otimes_a A),\ iA')$$

$$= \underline{Mor}_e(i\,sh(\Delta[n] \otimes_e iA),\ iA')$$

$$= \underline{Mor}_e(\Delta[n] \otimes_e iA,\ iA')$$

$$\underline{Hom}_e(\sharp A, A')_n = \underline{Mor}_e(\Delta[n] \otimes_e A,\ A')$$

$$= \underline{Mor}_a(sh(\Delta[n] \otimes_e iA),\ \sharp A')$$

$$= \underline{Mor}_e(\Delta[n] \otimes iA,\ iA') \qquad sh \dashv i$$

$$= \underline{Hom}_e(iA,\ iA')_n.$$

So a consequence of ~~def~~ definition $K \otimes_a A = sh(K \otimes_e iA)$

$$\therefore \underline{Hom}_a(A, A') = \underline{Hom}_e(iA, iA').$$

Now $\underline{Hom}_e(iA, i$

$$\underline{Mor}(K, \underline{Hom}_a(A, A')) = \underline{Mor}(K, \underline{Hom}_e(iA, iA')$$

$$= \underline{Hom}_e(iA,\ K \otimes iA')$$

Need right adjoint ~~exact~~ funct.

B. i: sheaves — ~~projective~~ presheaves

But sh: not a right ~~exact~~ adjoint of anything: does not preserve infinite products.

Cahiers Thomason. Bibliothèque MIR

Cahiers Thomason. Bibliothèque MIR

Cahiers Thomason. Bibliothèque MIR

RÉF. 801824 - 21 x 29,7 cm - Séyès
3 210338 018241

PARIS : 26 - 30 BOULEVARD St-Michel 6e
CLERMONT FERRAND - DIJON - GRENOBLE - LYON - MARSEILLE
MONTPELLIER - POITIERS - ST. ETIENNE - TOULOUSE - VERSAILLES

Cahiers Thomason, Bibliothèque MIR